

Nur für Forschungszwecke

NUDT8 Polyklonaler Antikörper

Katalog-Nr.:16098-1-AP



Allgemeine Informationen

Katalog-Nr.:
16098-1-AP

Größe:
150ul, Konzentration: 500 µg/ml von
Nanodrop und 347 µg/ml durch die
Bradford-Methode mit BSA als
Standard;

Wirt:
Kaninchen

Isotyp:
IgG

Immunogen Katalognummer:
AG8945

GenBank-Zugangsnummer:
BC018644

GeneID (NCBI):
254552

Vollständiger Name:
nudix (nucleoside diphosphate linked
moiety X)-type motif 8

Berechnete Masse:
236 aa, 25 kDa

Beobachtete Masse:
25 kDa

Reinigungsmethode:

Antigen-Affinitätsreinigung

Empfohlene Verdünnungen:

WB 1:1000-1:6000

IHC 1:50-1:500

Anwendungen

Geprüfte Anwendungen:

IHC, WB, ELISA

Getestete Reaktivität:

Human, Maus, Ratte

Hinweis-IHC: Antigendemaskierung mit TE-Puffer pH 9,0 empfohlen. (*) Wahlweise kann die Antigendemaskierung auch mit Citratpuffer pH 6,0 erfolgen.

Positivkontrollen:

WB: Jurkat-Zellen,

IHC: humanes Kolonkarzinomgewebe, humanes Eierstockgewebe, humanes Hautgewebe, humanes Herzgewebe, humanes Lebergewebe, humanes Milzgewebe, humanes Nierengewebe, Mausherzgewebe

Hintergrundinformationen

Lagerung

Lagerungsbedingungen:

Bei -20°C lagern. Nach dem Versand ein Jahr lang stabil

Lagerungspuffer:

PBS mit 0.02% Natriumazid und 50% Glycerin pH 7.3.

Aliquotieren ist nicht notwendig bei -20°C Lagerung

*** 20ul-Größen enthalten 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Ausgewählte Validierungsdaten




Jurkat cells were subjected to SDS PAGE followed by western blot with 16098-1-AP (NUDT8 antibody) at dilution of 1:1500 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded human heart using 16098-1-AP (NUDT8 antibody) at dilution of 1:100 (under 10x lens).




Immunohistochemical analysis of paraffin-embedded human heart using 16098-1-AP (NUDT8 antibody) at dilution of 1:100 (under 40x lens).




Jurkat cells were subjected to SDS PAGE followed by western blot with 16098-1-AP (NUDT8 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded mouse heart tissue slide using 16098-1-AP (NUDT8 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human colon cancer tissue slide using 16098-1-AP (NUDT8 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human colon cancer tissue slide using 16098-1-AP (NUDT8 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded mouse heart tissue slide using 16098-1-AP (NUDT8 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).