

# SMPDL3B Polyklonaler Antikörper

Katalog-Nr.: 16552-1-AP

Vorgestelltes Produkt

3 Publikationen

## Allgemeine Informationen

Katalog-Nr.:	16552-1-AP	GenBank-Zugangsnummer:	BC014444
Größe:	150ul, Konzentration: 600 µg/ml von Nanodrop und 333 µg/ml durch die Bradford-Methode mit BSA als Standard;	GenID (NCBI):	27293
Wirt:	Kaninchen	Vollständiger Name:	sphingomyelin phosphodiesterase, acid-like 3B
Isotyp:	IgG	Berechneté Masse:	373aa, 42 kDa; 455aa, 51 kDa
Immunogen Katalognummer:	AG9535	Beobachteté Masse:	42, 50-65 kDa

## Anwendungen

Geprüfte Anwendungen:	Positivkontrollen:
IHC, WB, ELISA	WB : COLO 320-Zellen, HEK-293-Zellen
In Publikationen genannte Anwendungen:	IHC : humanes Nierengewebe, humanes Kolonkarzinomgewebe
IHC, WB	
Getestete Reaktivität:	
Human, Maus, Ratte	
Zitierte Arten:	
Human, Maus, Ratte	
<b>Hinweis-IHC: Antigendemaskierung mit TE-Puffer pH 9,0 empfohlen. (*) Wahlweise kann die Antigendemaskierung auch mit Citratpuffer pH 6,0 erfolgen.</b>	

## Hintergrundinformationen

SMPDL3B is lipid-modulating phosphodiesterase. Active on the surface of macrophages and dendritic cells and strongly influences macrophage lipid composition and membrane fluidity. SMPDL3B acts as a negative regulator of Toll-like receptor signaling. It lacks activity with phosphocholine-containing lipids, but can cleave CDP-choline, and can release phosphate from ATP and ADP in vitro (PMID: 26095358). SMPDL3B can be detected 60-70 kDa by N-glycosylation modification (PMID: 26095358, PMID: 31217420).

## Bemerkenswerte Veröffentlichungen

Verfasser	Pubmed ID	Journal	Anwendung
Taesik Gwag	33294831	JHEP Rep	WB
Xin Li	29923306	Basic Clin Pharmacol Toxicol	WB,IHC
Huiqing Qu	34504869	Front Mol Biosci	WB

## Lagerung

Lagerungsbedingungen:  
Bei -20°C lagern. Nach dem Versand ein Jahr lang stabil  
Lagerungspuffer:  
PBS mit 0.02% Natriumazid und 50% Glycerin pH 7.3.  
Aliquotieren ist nicht notwendig bei -20°C Lagerung


\*\*\* 20ul-Größen enthalten 0.1% BSA

For technical support and original validation data for this product please contact:  
T: 1(888) 4PTGLAB (1-888-478-4522) (toll free  
in USA), or 1(312) 455-8498 (outside USA)


E: [proteintech@ptglab.com](mailto:proteintech@ptglab.com)  
W: [ptglab.com](http://ptglab.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Ausgewählte Validierungsdaten




COLO 320 cells lysates were subjected to SDS PAGE followed by western blot with 16552-1-AP (SMPDL3B antibody) at dilution of 1:1500 incubated at room temperature for 1.5 hours.




Various lysates were subjected to SDS PAGE followed by western blot with 16552-1-AP (SMPDL3B antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded human kidney tissue slide using 16552-1-AP (SMPDL3B antibody) at dilution of 1:500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human kidney tissue slide using 16552-1-AP (SMPDL3B antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human colon cancer tissue slide using 16552-1-AP (SMPDL3B antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human colon cancer tissue slide using 16552-1-AP (SMPDL3B antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).