

Nur für Forschungszwecke

# MYOD1 Polyklonaler Antikörper

Katalog-Nr.: 18943-1-AP

67 Publikationen



## Allgemeine Informationen

<b>Katalog-Nr.:</b> 18943-1-AP	<b>GenBank-Zugangsnummer:</b> BC064493	<b>Reinigungsmethode:</b> Antigen-Affinitätsreinigung
<b>Größe:</b> 150ul, Konzentration: 450 µg/ml von Nanodrop;	<b>GeneID (NCBI):</b> 4654	<b>Empfohlene Verdünnungen:</b> WB 1:1000-1:6000 IHC 1:50-1:500
<b>Wirt:</b> Kaninchen	<b>Vollständiger Name:</b> myogenic differentiation 1	
<b>Isotyp:</b> IgG	<b>Berechnete Masse:</b> 320 aa, 35 kDa	
<b>Immunogen Katalognummer:</b> AG13512	<b>Beobachtete Masse:</b> 35-45 kDa	

## Anwendungen

### Geprüfte Anwendungen:

FC, IHC, WB, ELISA

### In Publikationen genannte Anwendungen:

IF, IHC, WB

### Getestete Reaktivität:

Human, Maus, Ratte

### Zitierte Arten:

Ente, Hausschwein, Huhn, Human, Maus, Ratte, Ziege

**Hinweis-IHC: Antigendemaskierung mit TE-Puffer pH 9,0 empfohlen. (\*) Wahlweise kann die Antigendemaskierung auch mit Citratpuffer pH 6,0 erfolgen.**

### Positivkontrollen:

WB: Mausherzgewebe, humanes Herzgewebe, Maus-Skelettmuskelgewebe, Rattenherzgewebe

IHC: Mausembryogewebe,

## Hintergrundinformationen

MYOD1, also named as BHLHC1 or MYF3, is a 320 amino acid protein, which promotes the transcriptional activity of MYOD1 through its CDK9-mediated phosphorylation. This phosphorylation promotes its function in muscle differentiation. MYOD1 acts as a transcriptional activator that promotes transcription of muscle-specific target genes and plays a role in muscle differentiation. MYOD1 together with MYF5 and MYOG, co-occupies muscle-specific gene promoter core region during myogenesis. Induces fibroblasts to differentiate into myoblasts. The calculated molecular weight of MYOD1 is 34 kDa, but modified MYOD1 is about 45 kDa. (PMID: 12037670)

## Bemerkenswerte Veröffentlichungen

Verfasser	Pubmed ID	Journal	Anwendung
Yuanyuan Wu	36075558	Cell Signal	WB
Peng Ren	36118887	Front Genet	IHC
Hongyi Zhou	34502418	Int J Mol Sci	WB

## Lagerung

### Lagerungsbedingungen:

Bei -20°C lagern. Nach dem Versand ein Jahr lang stabil

### Lagerungspuffer:

PBS mit 0.02% Natriumazid und 50% Glycerin pH 7.3.

Aliquotieren ist nicht notwendig bei -20°C Lagerung

\*\*\* 20ul-Größen enthalten 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: [proteintech@ptglab.com](mailto:proteintech@ptglab.com)  
W: [ptglab.com](http://ptglab.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Ausgewählte Validierungsdaten



mouse heart tissue were subjected to SDS PAGE followed by western blot with 18943-1-AP (MYOD1 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded mouse embryo tissue slide using 18943-1-AP (MYOD1 Antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



$1 \times 10^6$  C2C12 cells were intracellularly stained with 0.4 ug Anti-Human MYOD1 (18943-1-AP) and CoraLite@488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).