

Nur für Forschungszwecke

APOB Polyklonaler Antikörper

Katalog-Nr.: 20578-1-AP

28 Publikationen



Allgemeine Informationen

Katalog-Nr.:	GenBank-Zugangsnummer:	Reinigungsmethode:
20578-1-AP	NM_000384	Antigen-Affinitätsreinigung
Größe:	GenID (NCBI):	Empfohlene Verdünnungen:
150ul, Konzentration: 500 µg/ml von Nanodrop;	338	WB 1:1000-1:8000 IHC 1:50-1:500 IF 1:20-1:200
Wirt:	Vollständiger Name:	
Kaninchen	apolipoprotein B (including Ag(x) antigen)	
Isotyp:	Berechneté Masse:	
IgG	516 kDa	
	Beobachteté Masse:	
	150,200-250 kDa, 400-520 kDa	

Anwendungen

Geprüfte Anwendungen:	Positivkontrollen:
IF, IHC, WB, ELISA	WB : human plasma, HEK-293-Zellen, humanes Plasmagewebe, Mauslebergewebe
In Publikationen genannte Anwendungen:	IHC : humanes Lebergewebe, humanes Leberzirrhosegewebe
IF, IHC, WB	IF : HepG2-Zellen,
Getestete Reaktivität:	
Human, Maus	
Zitierte Arten:	
Hamster, Human, Maus, Ratte	
Hinweis-IHC: Antigendemaskierung mit TE-Puffer pH 9,0 empfohlen. (*) Wahlweise kann die Antigendemaskierung auch mit Citratpuffer pH 6,0 erfolgen.	

Hintergrundinformationen

The apolipoprotein B (APOB) is a plasma protein synthesized primarily in the liver and intestine and play an important role in lipid and cholesterol metabolism. The APOB encodes two different isoproteins through mRNA editing, APOB-48 and APOB-100. APOB-48 and APOB-100 is present in both human liver and intestine. APOB-100 is essential for the assembly of VLDL in the liver. APOB-48 is essential for the assembly of chylomicrons in the intestine. It is well established that APOB-100 levels are associated with coronary heart disease. This antibody recognizes both of APOB-48, APOB-100 (PMID:11839763, PMID:2450346).

Bemerkenswerte Veröffentlichungen

Verfasser	Pubmed ID	Journal	Anwendung
Ying Zhang	36172518	Front Nutr	WB
Fangjun Yu	34493722	Nat Commun	WB
William A. Banks	36293369	Int J Mol Sci	WB

Lagerung

Lagerungsbedingungen:
Bei -20°C lagern. Nach dem Versand ein Jahr lang stabil
Lagerungspuffer:
PBS mit 0.02% Natriumazid und 50% Glycerin pH 7.3.
Aliquotieren ist nicht notwendig bei -20°C Lagerung


*** 20ul-Größen enthalten 0.1% BSA

For technical support and original validation data for this product please contact:
T: 1(888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Ausgewählte Validierungsdaten




Various lysates were subjected to Tris-acetate gel system followed by western blot with 20578-1-AP (APOB antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours.




mouse liver and human plasma were subjected to Tris-acetate gel system followed by western blot with 20578-1-AP (APOB antibody) at dilution of 1:1500 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded human liver tissue slide using 20578-1-AP (APOB antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffin-embedded human liver tissue slide using 20578-1-AP (APOB antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using 20578-1-AP (APOB antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunohistochemical analysis of paraffin-embedded human hepatocirrhosis tissue slide using 20578-1-AP (APOB antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human hepatocirrhosis tissue slide using 20578-1-AP (APOB antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).