

## Allgemeine Informationen

<b>Katalog-Nr.:</b> 22070-1-AP	<b>GenBank-Zugangsnummer:</b> BC000052	<b>Reinigungsmethode:</b> Antigen-Affinitätsreinigung
<b>Größe:</b> 150ul	<b>GeneID (NCBI):</b> 5465	
<b>Wirt:</b> Kaninchen	<b>Vollständiger Name:</b> peroxisome proliferator-activated receptor alpha	
<b>Isotyp:</b> IgG	<b>Berechnete Masse:</b> 468 aa, 52 kDa	
<b>Immunogen Katalognummer:</b> AG17192		

## Anwendungen

**Geprüfte Anwendungen:**  
ELISA

**Getestete Reaktivität:**  
Human

## Hintergrundinformationen

Peroxisome proliferator-activated receptor alpha (PPARA) is a ligand-activated transcription factor that belongs to the PPAR nuclear receptor superfamily. PPARA is essential in the modulation of lipid transport and metabolism, mainly through activating mitochondrial and peroxisomal fatty acid  $\beta$ -oxidation pathways. In addition, PPARA seems to decrease inflammation mainly through direct interaction with NF- $\kappa$ B, causing inhibition of its signaling pathway or reducing the activated levels of NF- $\kappa$ B and subsequent inflammation. Furthermore, PPARA was implicated in the attenuation of oxidative stress in alcoholic liver disease when treated with polyene phosphatidylcholine through downregulation of ROS-generating enzymes such as ethanol-inducible cytochrome P450 2E1 (CYP2E1), acyl-CoA oxidase, and NADPH oxidase. PPARA exists two isoforms and molecular weight of PPARA isoforms are 52 kDa and 22 kDa. The ability of a retinoid X receptor (RXR) to heterodimerize with many nuclear receptors, including LXR, PPAR, NGF1B and RAR, underscores its pivotal role within the nuclear receptor superfamily. Among these heterodimers, PPAR:RXR is considered an important signalling mediator of both PPAR ligands, such as fatty acids, and 9-cis retinoic acid (9-cis RA), an RXR ligand. (PMID: 15103326). PPARA can form Heterodimer with RXRA and molecular weight of Heterodimer is about 110 kDa.

## Lagerung

**Lagerungsbedingungen:**  
Bei -20°C lagern. Nach dem Versand ein Jahr lang stabil

**Lagerungspuffer:**  
PBS mit 0.02% Natriumazid und 50% Glycerin pH 7.3.

**Aliquotieren ist nicht notwendig bei -20°C Lagerung**

**\*\*\* 20ul-Größen enthalten 0.1% BSA**

