

Nur für Forschungszwecke

MMP9 (Middle) Polyklonaler Antikörper



Katalog-Nr.: 27306-1-AP

20 Publikationen

Allgemeine Informationen

Katalog-Nr.:
27306-1-AP

Größe:
150ul, Konzentration: 350 µg/ml von
Nanodrop und 267 µg/ml durch die
Bradford-Methode mit BSA als
Standard;

Wirt:
Kaninchen

Isotyp:
IgG

Immunogen Katalognummer:
AG26132

GenBank-Zugangsnummer:
BC006093

GeneID (NCBI):
4318

Vollständiger Name:
matrix metalloproteinase 9
(gelatinase B, 92kDa gelatinase,
92kDa type IV collagenase)

Berechnete Masse:
707 aa, 78 kDa

Beobachtete Masse:
92 kDa

Reinigungsmethode:

Antigen-Affinitätsreinigung

Empfohlene Verdünnungen:

WB 1:500-1:1000
IP 0.5-4.0 µg für IP und 1:200-1:1000
für WB
IHC 1:50-1:500

Anwendungen

Geprüfte Anwendungen:

IHC, IP, WB, ELISA

In Publikationen genannte Anwendungen:

IHC, WB

Getestete Reaktivität:

Human

Zitierte Arten:

Human, Maus, Ratte

Hinweis-IHC: Antigendemaskierung mit TE-Puffer pH 9,0 empfohlen. (*) Wahlweise kann die Antigendemaskierung auch mit Citratpuffer pH 6,0 erfolgen.

Positivkontrollen:

WB: humanes Speichelgewebe,

IP: humanes Speichelgewebe,

IHC: humanes Tonsillitisgewebe, humanes Milzgewebe

Hintergrundinformationen

Proteins of the matrix metalloproteinase (MMP) family are involved in the breakdown of extracellular matrix in normal physiological processes, such as embryonic development, reproduction, tissue remodeling, and disease processes, such as arthritis or metastasis. Most MMP's are secreted as inactive proproteins which are activated when cleaved by extracellular proteinases. Matrix metalloproteinase 9 (gelatinase B, 92kDa gelatinase, 92kDa type IV collagenase) (MMP9, synonyms: GELB, CLG4B) degrades collagens type IV and V. Studies in rhesus monkeys suggest that MMP9 is involved in IL-8-induced mobilization hematopoietic progenitor cells from bone marrow, and murine studies suggest a role in tumor-associated tissue remodeling. The pro-MMP9 is 92 kDa, and it can be detected a processed form of 68 kDa or 82 kDa. This protein can exist as a dimer of 180 kDa (PMID:7492685).

Bemerkenswerte Veröffentlichungen

Verfasser	Pubmed ID	Journal	Anwendung
XIAOYUE FENG	34528694	Oncol Rep	WB
WANG Xiao-He	34688464	Chin J Nat Med	WB
Cong Xu	34868365	Oncol Lett	WB

Lagerung

Lagerungsbedingungen:

Bei -20°C lagern. Nach dem Versand ein Jahr lang stabil

Lagerungspuffer:

PBS mit 0.02% Natriumazid und 50% Glycerin pH 7.3.

Aliquotieren ist nicht notwendig bei -20°C Lagerung

*** 20ul-Größen enthalten 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Ausgewählte Validierungsdaten




human saliva were subjected to SDS PAGE followed by western blot with 27306-1-AP (MMP9 (Middle) antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.



IP result of anti-MMP9 (Middle)(IP:27306-1-AP, 4ug; Detection:27306-1-AP 1:300) with human saliva lysate 800 ug.



Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 27306-1-AP (MMP9 (Middle) antibody) at dilution of 1:200 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 27306-1-AP (MMP9 (Middle) antibody) at dilution of 1:200 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).