

Nur für Forschungszwecke

CHD8 Polyklonaler Antikörper

Katalog-Nr.:29783-1-AP



Allgemeine Informationen

Katalog-Nr.: 29783-1-AP	GenBank-Zugangsnummer: NM_001170629	Reinigungsmethode: Antigen-Affinitätsreinigung
Größe: 150ul , Konzentration: 750 µg/ml von Nanodrop;	GeneID (NCBI): 57680	Empfohlene Verdünnungen: WB 1:2000-1:12000 IHC 1:250-1:1000
Wirt: Kaninchen	Vollständiger Name: chromodomain helicase DNA binding protein 8	
Isotyp: IgG	Berechnete Masse: 290KD	
Immunogen Katalognummer: AG31425	Beobachtete Masse: 290 kDa	

Anwendungen

Geprüfte Anwendungen:
IHC, WB, ELISA

Getestete Reaktivität:
Human, Maus, Ratte

Hinweis-IHC: Antigenmaskierung mit TE-Puffer pH 9,0 empfohlen. (*) Wahlweise kann die Antigenmaskierung auch mit Citratpuffer pH 6,0 erfolgen.

Positivkontrollen:

WB : HeLa-Zellen, Maushirngewebe, Rattenhirngewebe

IHC : Maushirngewebe,

Hintergrundinformationen

Chromodomain helicase domain 8 (CHD8) is one of the most frequently mutated and most penetrant genes in the autism spectrum disorder (ASD). CHD8 is located on 14q11.2. It is part of the SNF2H-like ATP-dependent chromatin remodeling enzymes family referred to as CHD (chromodomain helicase DNA binding). CHD8 has two isoforms: CHD8L, a full-length protein of 280 kDa; and CHD8S (Duplin), a 110 kDa protein of the NH2-terminal chromodomain region resulting from alternative splicing (PMID:19151705). CHD8 is essential for development, as homozygote mutant mice die at an embryonic stage. CHD8 is expressed in the mouse at the embryonic stage (E12.5) in different levels in wide regions of the brain (neocortex, forebrain, ventricular, subventricular and mantle zones, rhombic lip (RL), and the isthmus of the cerebellum, as well as in lower RL and floor plate region of the hindbrain, midbrain, diencephalon, hypothalamus, pituitary gland, craniofacial region, and tongue and olfactory epithelium). In the postnatal mouse brain (P20), CHD8 is expressed in the cerebellum, neocortex, hippocampus, hypothalamus, and olfactory bulb (PMID:30277262). Peak expression levels were observed at E18-P7, then gradually decreased to adulthood. Highest expression was found in neurons, and lower levels in astrocyte and astroglia (PMID:30574290). In the mouse, CHD8 is expressed higher in brain compared to other tissues and in the embryo, compared to adult.

Lagerung

Lagerungsbedingungen:

Bei -20°C lagern.

Lagerungspuffer:

PBS mit 0.02% Natriumazid und 50% Glycerin pH 7.3.

Aliquotieren ist nicht notwendig bei -20°C Lagerung

*** 20ul-Größen enthalten 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Ausgewählte Validierungsdaten



Various lysates were subjected to SDS PAGE followed by western blot with 29783-1-AP (CHD8 antibody) at dilution of 1:6000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded mouse brain tissue slide using 29783-1-AP (CHD8 antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded mouse brain tissue slide using 29783-1-AP (CHD8 antibody) at dilution of 1:500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).