

Nur für Forschungszwecke

# SCAMP5 Polyklonaler Antikörper

Katalog-Nr.:30328-1-AP



## Allgemeine Informationen

<b>Katalog-Nr.:</b> 30328-1-AP	<b>GenBank-Zugangsnummer:</b> BC024700	<b>Reinigungsmethode:</b> Antigen-Affinitätsreinigung
<b>Größe:</b> 150ul , Konzentration: 600 µg/ml von Nanodrop;	<b>GeneID (NCBI):</b> 192683	<b>Empfohlene Verdünnungen:</b> WB 1:1000-1:8000
<b>Wirt:</b> Kaninchen	<b>Vollständiger Name:</b> secretory carrier membrane protein 5	
<b>Isotyp:</b> IgG	<b>Berechnete Masse:</b> 26 kDa	
<b>Immunogen Katalognummer:</b> AG31837	<b>Beobachtete Masse:</b> 26 kDa	

## Anwendungen

<b>Geprüfte Anwendungen:</b> WB, ELISA	<b>Positivkontrollen:</b> WB : Maushirngewebe, mouse brain tissue (incubated at 37°C)
<b>Getestete Reaktivität:</b> Human, Maus	

## Hintergrundinformationen

Secretory carrier-associated membrane protein 5 (SCAMP5) belongs to the SCAMP family. Secretory carrier membrane proteins (SCAMPs) constitute a group of membrane transport proteins in plants, insects and mammals. The mammalian genome contains five types of SCAMP genes, namely, SCAMP1-SCAMP5. (PMID:36217917, PMID:19234194). SCAMPs participate in the vesicle cycling fusion of vesicles and cell membranes and play roles in regulating exocytosis and endocytosis, activating synaptic function and transmitting nerve signals. Among these proteins, SCAMP5 is highly expressed in the brain and has direct or indirect effects on the function of the central nervous system (PMID:36217917). SCAMP5 regulates membrane transport, controls the exocytosis of synaptic vesicles and is related to secretion carrier and membrane function. In addition, SCAMP5 plays a major role in the normal maintenance of the physiological functions of nerve cells (PMID:36217917, PMID:33663553). For optimal WB detection of this membrane protein, we recommend to avoid boiling the sample after lysis.

## Lagerung


**Lagerungsbedingungen:**  
Bei -20°C lagern. Nach dem Versand ein Jahr lang stabil  
**Lagerungspuffer:**  
PBS mit 0.02% Natriumazid und 50% Glycerin pH 7.3.  
**Aliquotieren ist nicht notwendig bei -20°C Lagerung**

**\*\*\* 20ul-Größen enthalten 0.1% BSA**

For technical support and original validation data for this product please contact:  
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)  
E: proteintech@ptglab.com  
W: ptglab.com

**This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.**

## Ausgewählte Validierungsdaten



Various lysates were subjected to SDS PAGE followed by western blot with 30328-1-AP (SCAMP5 antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours.