

Nur für Forschungszwecke

# ADRP/Perilipin 2 Monoklonaler Antikörper

Katalog-Nr.: **60340-1-Ig**



## Allgemeine Informationen

Katalog-Nr.:	60340-1-Ig	GenBank-Zugangsnummer:	BC005127	Reinigungsmethode:	Protein-G-Reinigung
Größe:	150ul , Konzentration: 1900 µg/ml von 123 Nanodrop und 1167 µg/ml durch die Bradford-Methode mit BSA als Standard;	GenID (NCBI):	Vollständiger Name: adipose differentiation-related protein	CloneNo.:	1G5D4
Wirt:	Maus	Berechneté Masse:	48 kDa	Empfohlene Verdünnungen:	WB 1:500-1:2000 IHC 1:200-1:8000
Isotyp:	IgG1	Beobachteté Masse:	47-52 kDa		
Immunogen Katalognummer:	AG7643				

## Anwendungen

Geprüfte Anwendungen:	Positivkontrollen:
IHC, WB, ELISA	WB : humanes Fettgewebe,
Getestete Reaktivität:	IHC : humanes Nierenzellkarzinomgewebe, braunes Mausfettgewebe
<b>Hinweis-IHC: Antigendemaskierung mit TE-Puffer pH 9,0 empfohlen. (*) Wahlweise kann die Antigendemaskierung auch mit Citratpuffer pH 6,0 erfolgen.</b>	

## Hintergrundinformationen

ADRP (adipocyte differentiation related protein) also known as ADFP, adipophilin, or perilipin-2, is a member of PAT family which is responsible for the transportation of lipids and the formation of lipid droplets. ADRP is localized on the surface of lipid droplets in a variety of tissues and cell lines. ADRP is not detected in undifferentiated cells but increases rapidly to high levels when adipocyte precursors differentiate into adipocytes. Anti-ADRP antibody is a reliable and sensitive marker for lipid droplet. Enhanced expression of ADRP is linked to diseases with abnormal lipid storage, including hepatic steatosis, atherosclerosis and diabetes. Immunohistochemistry of ADRP may facilitate histomorphological diagnosis of these diseases.

## Lagerung

Lagerungsbedingungen:  
Bei -20°C lagern. Nach dem Versand ein Jahr lang stabil  
Lagerungspuffer:  
PBS mit 0.02% Natriumazid und 50% Glycerin pH 7.3.  
Aliquotieren ist nicht notwendig bei -20°C Lagerung




\*\*\* 20ul-Größen enthalten 0.1% BSA

For technical support and original validation data for this product please contact:  
T: 1(888) 4PTGLAB (1-888-478-4522) (toll free  
in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com  
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Ausgewählte Validierungsdaten




human adipose tissue were subjected to SDS PAGE followed by western blot with 60340-1-Ig (ADRP Antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.

Immunohistochemical analysis of paraffin-embedded human renal cell carcinoma tissue slide using 60340-1-Ig (ADRP Antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).


Immunohistochemical analysis of paraffin-embedded human renal cell carcinoma tissue slide using 60340-1-Ig (ADRP Antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded rat brown adipose slide using 60340-1-Ig (ADRP/Perilipin 2 antibody) at dilution of 1:16000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human renal cell carcinoma tissue slide using 60340-1-Ig (ADRP/Perilipin 2 antibody) at dilution of 1:8000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human renal cell carcinoma tissue slide using 60340-1-Ig (ADRP/Perilipin 2 antibody) at dilution of 1:8000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).