Nur für Forschungszwecke

CD16 Monoklonaler Antikörper

Katalog-Nr.:65090-1-lg





Allgemeine Informationen	Katalog-Nr.: 65090-1-lg	GenBank-Zugangsnummer: BC017865	Reinigungsmethode: Affinitätsreinigung
	Größe: 100ug , 0.5 mg/ml	GenelD (NCBI): 2214	CloneNo.: 3G8
	Wirt: Maus Isotyp: IgG1, kappa	Vollständiger Name: Fc fragment of IgG, low affinity III receptor (CD16a)	a,
		Berechneté Masse: 254 aa, 29 kDa	
Anwendungen	Geprüfte Anwendungen: FC Getestete Reaktivität: Human		
Hintergrundinformationen	CD16 is a 50-70-kDa low affinity Fc receptor found on the surface of natural killer cells, neutrophil polymorphonuclear leukocytes, monocytes and macrophages. CD16 mediates antibody-dependent cellular cytotoxicity (ADCC) and other antibody-dependent responses, such as phagocytosis. CD16 has been identified as Fc receptors Fcy RIIIa (CD16a) and Fcy RIIIb (CD16b), encoded by two nearly identical genes, FCGR3A and the FCGR3B. Clone 3G8 recognizes both the CD16a and CD16b (PMID: 7592758).		
Lagerung	Lagerungsbedingungen: Store at 2-8°C. Stable for one y Lagerungspuffer: PBS with 0.09% sodium azide.	ear after shipment.	

For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll freeE: proteintech@ptglab.comin USA), or 1(312) 455-8498 (outside USA)W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Ausgewählte Validierungsdaten





1X10⁶ human PBMCs were surface co-stained with FITC Anti-Human CD3 and 0.2 ug Anti-Human CD16 (65090-1-Ig, Clone:3G8) and APC-Goat anti-Mouse IgG at dilution 1:1000. Cells were not fixed. Lymphocytes were gated.

1X10^{^6} human PBMCs were surface stained with 0.2 ug Anti-Human CD16 (65090-1-1g, Clone:3G8) and APC-Goat anti-Mouse IgG at dilution 1:1000. Cells were not fixed. Lymphocytes were gated.