Nur für Forschungszwecke

CD357 (GITR) Monoklonaler Antikörper

Katalog-Nr.:65102-1-Ig




Allgemeine Informationen	Katalog-Nr.: 65102-1-lg	GenBank-Zugangsnummer: BC146517	Reinigungsmethode: Affinitätsreinigung
	<mark>Größe:</mark> 100ug , 500 µg/ml	GenelD (NCBI): 21936	CloneNo.: DTA-1
	Wirt: Ratte	Vollständiger Name: tumor necrosis factor receptor superfamily, member 18	
	<mark>Isotyp:</mark> IgG2b, kappa		
Anwendungen	Geprüfte Anwendungen: FC		
	Getestete Reaktivität: Maus		
Hintergrundinformationen	Glucocorticoid-induced TNFR-related protein (GITR), also known as CD357 or TNFRSF18, is a member of the tumor necrosis factor receptor (TNF-R) superfamily. GITR is expressed constitutively at high levels in T regulatory cells (Treg cells) and plays a key role in dominant immunological self-tolerance maintained by CD25+CD4+ regulatory T cells (PMID: 11812990). It is expressed at low levels on resting responder T cells. The expression of GITR on T cells can be upregulated upon activation (PMID: 15770698). GITR is activated by GITR ligand (GITRL) which is mainly expressed on APC. GITR-GITRL interactions could co-stimulate both responder T-cell functions and the suppressive functions of Treg cells (PMID: 16868552).		
Lagerung	Lagerungsbedingungen: Store at 2-8°C. Stable for one year at Lagerungspuffer: PBS mit 0,1% Natriumazid, pH 7,3.	fter shipment.	

For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll freeE: proteintech@ptglab.comin USA), or 1(312) 455-8498 (outside USA)W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Ausgewählte Validierungsdaten



1X10^6 mouse splenocytes were surface stained with PE-Anti-Mouse CD4 (PE-65104, Clone: GK1.5), CoraLite® Plus 647-Conjugated Anti-Mouse CD25 (CL647-65137, Clone: PC61.5), 0.5 ug Anti-Mouse CD357 (GITR) (65102-1-1g, Clone: DTA-1) and FITC anti-rat IgG2b antibody at dilution 1:200. Cells were not fixed. CD4+ lymphocytes were gated for analysis of CD25 and CD357 (GITR) staining.