

Nur für Forschungszwecke

MHC Class II (I-A/I-E) Monoklonaler Antikörper



Katalog-Nr.: 65122-1-Ig **1 Publikationen**

Allgemeine Informationen

Katalog-Nr.: 65122-1-Ig	GenBank-Zugangsnummer: BC031711	Reinigungsmethode: Affinitätsreinigung
Größe: 100ug, 0.5 mg/ml	GeneID (NCBI): 14960	CloneNo.: M5/114.15.2
Wirt: Ratte	Vollständiger Name: histocompatibility 2, class II antigen	
Isotyp: IgG2b, kappa	A, alpha	

Anwendungen

Geprüfte Anwendungen:
FC

In Publikationen genannte Anwendungen:
IF

Getestete Reaktivität:
Maus

Zitierte Arten:
Maus

Hintergrundinformationen

Major histocompatibility complex (MHC) class II molecules are heterodimeric transmembrane glycoproteins that are expressed on the surface of antigen-presenting cells such as dendritic cells, macrophages, and B cells. The M5/114.15.2 antibody detects polymorphic determinants present on B cells, monocytes, macrophages, dendritic cells, and activated T lymphocytes from mice carrying the H-2b, H-2d, H-2q, H-2p, H-2r and H-2u haplotypes but not from mice carrying the H-2s or H-2f haplotypes (PMID: 6170707). This antibody recognizes both I-A and I-E subregion-encoded Ia glycoproteins (I-Ab, I-Ad, I-Aq, I-Ed, I-Ek, not I-Af, I-Ak, or I-As). M5/114 mAb is reported to inhibit I-A subregion-restricted T cell responses of the H-2b, H-2d, H-2q, H-2u but not the H-2f, H-2k, H-2s haplotypes (PMID: 6173436).

Bemerkenswerte Veröffentlichungen

Verfasser	Pubmed ID	Journal	Anwendung
Yuwei Zhang	35219165	Int Immunopharmacol	IF

Lagerung


Lagerungsbedingungen:
Store at 2-8°C. Stable for one year after shipment.

Lagerungspuffer:
PBS with 0.09% sodium azide.

For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Ausgewählte Validierungsdaten



1X10⁶ mouse splenocytes were surface stained with 0.5 µg Anti-Mouse MHC Class II (I-A/I-E) (65122-1-Ig, Clone: M5/114.15.2) and FITC anti-rat IgG2b Antibody at dilution 1:100 (red), or 0.5 µg Isotype Control. Cells were not fixed.