

## Allgemeine Informationen

**Katalog-Nr.:**

65168-1-Ig

**Größe:**

100 µg, 0.5 mg/ml

**Wirt:**

Maus

**Isotyp:**

IgG1, kappa

**GenBank-Zugangsnummer:**

BC074740

**GeneID (NCBI):**

5133

**Vollständiger Name:**

programmed cell death 1

**Berechnete Masse:**

288 aa, 32 kDa

**Reinigungsmethode:**

Mit Protein-A-

Affinitätschromatographie gereinigt

**CloneNo.:**

EH12.2H7

## Anwendungen

**Geprüfte Anwendungen:**

FC

**Getestete Reaktivität:**

Human, Nicht-Humane Primaten

## Hintergrundinformationen

Programmed cell death 1 (PD-1, also known as CD279) is an immunoinhibitory receptor that belongs to the CD28/CTLA-4 subfamily of the Ig superfamily. It is a 288 amino acid (aa) type I transmembrane protein composed of one Ig superfamily domain, a stalk, a transmembrane domain, and an intracellular domain containing an immunoreceptor tyrosine-based inhibitory motif (ITIM) as well as an immunoreceptor tyrosine-based switch motif (ITSM) (PMID: 18173375). PD-1 is expressed during thymic development and is induced in a variety of hematopoietic cells in the periphery by antigen receptor signaling and cytokines (PMID: 20636820). Engagement of PD-1 by its ligands PD-L1 or PD-L2 transduces a signal that inhibits T-cell proliferation, cytokine production, and cytolytic function (PMID: 19426218). It is critical for the regulation of T cell function during immunity and tolerance. Blockade of PD-1 can overcome immune resistance and also has been shown to have antitumor activity (PMID: 22658127; 23169436).

## Lagerung

**Lagerungsbedingungen:**

Store at 2-8°C. Stable for one year after shipment.

**Lagerungspuffer:**

PBS with 0.09% sodium azide.


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: [proteintech@ptglab.com](mailto:proteintech@ptglab.com)  
W: [ptglab.com](http://ptglab.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Ausgewählte Validierungsdaten



$1 \times 10^6$  day 3 PHA treated human PBMCs were surface stained with 0.2  $\mu$ g Anti-Human CD279 (65168-1-Ig, Clone:EH12.2H7) and CL594-Goat anti-Mouse IgG at dilution 1:1000. Cells were not fixed. Lymphocytes were gated. Blue line indicates unstained stimulated cells. Black dashed line indicates unstimulated stained cells.