Nur für Forschungszwecke

ALCAM Monoklonaler Antikörper

Katalog-Nr.:65223-1-Ig



Reinigungsmethode:

Affinitätsreinigung

CloneNo.:

3A6

Allgemeine Informationen

Katalog-Nr.: 65223-1-lg

Größe:

100ug , 0.5 mg/mL

Wirt: Maus

Isotyp:

IgG1, kappa

105 kDa

molecule

BC057809

GeneID (NCBI):

Vollständiger Name:

Berechneté Masse:

GenBank-Zugangsnummer:

activated leukocyte cell adhesion

Geprüfte Anwendungen:

Getestete Reaktivität:

Human

Hintergrundinformationen

Activated leukocyte cell adhesion molecule (ALCAM, also known as CD166) is a cell adhesion molecule that belongs to the immunoglobulin superfamily. It is involved in cell-cell adhesion through homophilic and heterophilic (to CD6) interactions. ALCAM is widely expressed in a variety of normal tissues and cell types, including activated T cells and monocytes, epithelial cells, fibroblasts, neuronal cells, hepatocytes, and bone marrow mesenchymal stem cells (PMID: 7760007; 25221999). Altered ALCAM expression has been associated with $the \ differentiation \ state \ and \ progression \ in \ some \ neoplasms \ including \ melanoma, \ prostate, \ colorectal, \ and \ breast$ cancers (PMID: 20461761; 18172759).

Lagerung

Anwendungen

Lagerungsbedingungen:

Store at 2-8°C. Stable for one year after shipment.

Lagerungspuffer:


PBS with 0.09% sodium azide.

in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Ausgewählte Validierungsdaten



1X10^6 PHA treated human PBMCs were surface stained with 0.2 ug Anti-Human ALCAM (65223-1-Ig, Clone:3A6) and CL594-Goat anti-Mouse IgG at dilution 1:1000 (red) or unstained (blue). Cells were not fixed. Monocytes were gated.