

CD72 Monoklonaler Antikörper

Katalog-Nr.:**67424-1-Ig**

Allgemeine Informationen

Katalog-Nr.:	GenBank-Zugangsnummer:
67424-1-Ig	BC030227
Größe:	GenID (NCBI):
150ul , Konzentration: 2200 µg/ml von 971 Nanodrop und 1000 µg/ml durch die Bradford-Methode mit BSA als Standard;	Vollständiger Name: CD72 molecule
Wirt:	Berechneté Masse: 359 aa, 40 kDa
Istotyp:	Beobachteté Masse: 40 kDa
Immunogen Katalognummer:	
AG14405	

Anwendungen

Geprüfte Anwendungen:	Positivkontrollen:
FC, IF, IHC, WB, ELISA	WB : Daudi-Zellen,
Getestete Reaktivität:	IHC : humanes Tonsillitisgewebe,
Human	IF : humanes Tonsillitisgewebe,

Hinweis-IHC: Antigendemaskierung mit TE-Puffer pH 9,0 empfohlen. (*) Wahlweise kann die Antigendemaskierung auch mit Citratpuffer pH 6,0 erfolgen.

Hintergrundinformationen

CD72 is a type II transmembrane glycoprotein of 39-45 kDa and is expressed as a disulphide-linked homodimer (PMID: 1711157). CD72 is a pan B-cell marker covering the full range of differentiation from the very early stages of B cells to mature surface Ig+ B cells (PMID: 1384316). It is expressed in all stages of B cell development except plasma cells. CD72 is also expressed by dendritic cells, tissue macrophages in the red pulp of the spleen and von Kupffer cells in the liver (PMID: 8561576; 1384316). CD72 is the ligand for CD5 (PMID: 1711157). CD72 is a negative regulator of BCR signaling (PMID: 9590210). Interaction of CD100 with CD72 strictly tunes the strength of BCR signals and is essential for maintaining immunological homeostasis as well as generating a proper immune response (PMID: 16113236).

Lagerung





Lagerungsbedingungen:
Bei -20°C lagern.
Lagerungspuffer:
PBS mit 0.02% Natriumazid und 50% Glycerin pH 7.3.
Aliquotieren ist nicht notwendig bei -20°C Lagerung

*** 20ul-Größen enthalten 0.1% BSA

Ausgewählte Validierungsdaten



Daudi cells were subjected to SDS PAGE followed by western blot with 67424-1-Ig (CD72 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



1X10⁶ Ramos cells were stained with 0.20 ug Anti-Human CD72 (67424-1-Ig, Clone:2C1C6) (red) or 0.20 ug control antibody (blue) and CoraLite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) with dilution of 1:1000. Cells were fixed with 90% MeOH.