

Nur für Forschungszwecke

TCF7L1/TCF7L2 Monoklonaler Antikörper

Katalog-Nr.:67652-1-Ig



Allgemeine Informationen

Katalog-Nr.: 67652-1-Ig	GenBank-Zugangsnummer: BC058894	Reinigungsmethode: Protein-A-Reinigung
Größe: 150ul , Konzentration: 2100 µg/ml von83439 Nanodrop und 1000 µg/ml durch die Bradford-Methode mit BSA als Standard;	GeneID (NCBI): transcription factor 7-like 1 (T-cell specific, HMG-box)	CloneNo.: 2B2D12
Wirt: Maus	Berechnete Masse: 63 kDa	Empfohlene Verdünnungen: WB 1:20000-1:100000
Isotyp: IgG2b	Beobachtete Masse: 70 kDa	
Immunogen Katalognummer: AG28954		

Anwendungen

Geprüfte Anwendungen: WB,ELISA	Positivkontrollen: WB : HEK-293-Zellen, A431-Zellen, HeLa-Zellen, JAR-Zellen, NCCIT-Zellen, NIH/3T3-Zellen
Getestete Reaktivität: Human, Maus	

Hintergrundinformationen

TCF7L1 is one of TCF/LEF transcription factors that are mediators of the Wnt signaling pathway and are antagonized by the TGF-beta signaling pathway. It binds to DNA and acts as a repressor in the absence of CTNNB1, and as an activator in its presence. It's necessary for the terminal differentiation of epidermal cells, the formation of keratohyalin granules and the development of the barrier function of the epidermis. This antibody raised against 1-200aa of human TCF7L1 can also react with TCF7L2.

Lagerung


Lagerungsbedingungen:
Bei -20°C lagern.
Lagerungspuffer:
PBS mit 0.02% Natriumazid und 50% Glycerin pH 7.3.
Aliquotieren ist nicht notwendig bei -20°C Lagerung

***** 20ul-Größen enthalten 0.1% BSA**

For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Ausgewählte Validierungsdaten



Various lysates were subjected to SDS PAGE followed by western blot with 67652-1-Ig (TCF7L1/TCF7L2 antibody) at dilution of 1:50000 incubated at room temperature for 1.5 hours.