

## Allgemeine Informationen

<b>Katalog-Nr.:</b> 67742-1-Ig	<b>GenBank-Zugangsnummer:</b> BC005377	<b>Reinigungsmethode:</b> Protein-G-Reinigung
<b>Größe:</b> 150ul , Konzentration: 1000 µg/ml von34 Nanodrop und 449 µg/ml durch die Bradford-Methode mit BSA als Standard;	<b>GeneID (NCBI):</b> 1A11G6	<b>CloneNo.:</b> 1A11G6
<b>Wirt:</b> Maus	<b>Vollständiger Name:</b> acyl-Coenzyme A dehydrogenase, C-4 to C-12 straight chain	<b>Empfohlene Verdünnungen:</b> IWB 1:2000-1:10000 IHC 1:250-1:1000 IF 1:200-1:800
<b>Isotyp:</b> IgG1	<b>Berechnete Masse:</b> 421 aa, 47 kDa	
<b>Immunogen Katalognummer:</b> AG30620	<b>Beobachtete Masse:</b> 42 kDa	

## Anwendungen

### Geprüfte Anwendungen:

IF, IHC, WB, ELISA

### Getestete Reaktivität:

Hausschwein, Human, Ratte

**Hinweis-IHC: Antigendemaskierung mit TE-Puffer pH 9,0 empfohlen. (\*) Wahlweise kann die Antigendemaskierung auch mit Citratpuffer pH 6,0 erfolgen.**

### Positivkontrollen:

**WB:** PC-3-Zellen, HEK-293-Zellen, HeLa-Zellen, HepG2-Zellen, Rattenlebergewebe

**IHC:** humanes Lebergewebe, humanes Herzgewebe

**IF:** humanes Leberkarzinomgewebe,

## Hintergrundinformationen

ACADM, also named as MCAD, belongs to the acyl-CoA dehydrogenase family. This enzyme is specific for acyl chain lengths of 4 to 16. It catalyzes the reaction: Acyl-CoA + acceptor = 2,3-dehydroacyl-CoA + reduced acceptor. Defects in ACADM are the cause of medium-chain acyl-CoA dehydrogenase deficiency (MCAD deficiency). This protein can exist as a dimer(PMID:8962055). This antibody is specific to ACADM.

## Lagerung

### Lagerungsbedingungen:

Bei -20°C lagern. Nach dem Versand ein Jahr lang stabil

### Lagerungspuffer:

PBS mit 0.02% Natriumazid und 50% Glycerin pH 7.3.

Aliquotieren ist nicht notwendig bei -20°C Lagerung

\*\*\* 20ul-Größen enthalten 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com  
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Ausgewählte Validierungsdaten




Various lysates were subjected to SDS PAGE followed by western blot with 67742-1-Ig (ACADM antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.




rat liver tissue were subjected to SDS PAGE followed by western blot with 67742-1-Ig (ACADM antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded human liver tissue slide using 67742-1-Ig (ACADM antibody) at dilution of 1:500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human liver tissue slide using 67742-1-Ig (ACADM antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed human liver cancer tissue using ACADM antibody (67742-1-Ig, Clone: 1A11G6) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).



HSC-T6 cells were subjected to SDS PAGE followed by western blot with 67742-1-Ig (ACADM antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.