

Nur für Forschungszwecke

# PLEKHB2 Monoklonaler Antikörper

Katalog-Nr.:67877-1-Ig



## Allgemeine Informationen

<b>Katalog-Nr.:</b> 67877-1-Ig	<b>GenBank-Zugangsnummer:</b> BC013991	<b>Reinigungsmethode:</b> Protein-G-Reinigung
<b>Größe:</b> 150ul, Konzentration: 1000 µg/ml von55041 Nanodrop;	<b>GeneID (NCBI):</b> 55041	<b>CloneNo.:</b> 1H9D7
<b>Wirt:</b> Maus	<b>Vollständiger Name:</b> pleckstrin homology domain containing, family B (evectins) member 2	<b>Empfohlene Verdünnungen:</b> WB 1:2000-1:10000
<b>Isotyp:</b> IgG1	<b>Berechnete Masse:</b> 222 aa, 25 kDa	
<b>Immunogen Katalognummer:</b> AG9231	<b>Beobachtete Masse:</b> 19-25;30-34 kDa	

## Anwendungen

<b>Geprüfte Anwendungen:</b> WB, ELISA	<b>Positivkontrollen:</b> WB : Caco-2-Zellen, BxPC-3-Zellen, Hausschwein-Kolongewebe, HL-60-Zellen, HT-29-Zellen, NCCIT-Zellen, SW480-Zellen, U-87 MG-Zellen
<b>Getestete Reaktivität:</b> Human	

## Hintergrundinformationen

PLEKHB2, also named as EVT2 (Evectin-2), is involved in retrograde transport of recycling endosomes. PLEKHB2 has been identified as post-Golgi proteins of unknown function that have a PH domain that typically binds polyphosphoinositides. PLEKHB2 is expressed in a broad range of tissues. PLEKHB2 has some isoforms with MW 19-26 kDa and 34 kDa. This antibody recognizes all the isoforms of PLEKHB2.

## Lagerung

**Lagerungsbedingungen:**  
Bei -20°C lagern. Nach dem Versand ein Jahr lang stabil  
**Lagerungspuffer:**  
PBS mit 0.02% Natriumazid und 50% Glycerin pH 7.3.  
Aliquotieren ist nicht notwendig bei -20°C Lagerung

\*\*\* 20ul-Größen enthalten 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com  
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Ausgewählte Validierungsdaten



Various lysates were subjected to SDS PAGE followed by western blot with 67877-1-Ig (PLEKHB2 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated GAPDH Monoclonal antibody (HRP-60004) as loading control.