

Nur für Forschungszwecke

TOLLIP Monoklonaler Antikörper

Katalog-Nr.:68170-1-Ig



Allgemeine Informationen

Katalog-Nr.: 68170-1-Ig	GenBank-Zugangsnummer: BC012057	Reinigungsmethode: Protein-G-Reinigung
Größe: 150ul , Konzentration: 1000 µg/ml von54472	GeneID (NCBI): 54472	CloneNo.: 1B3C4
Nanodrop;	Vollständiger Name: toll interacting protein	Empfohlene Verdünnungen: WB 1:5000-1:50000
Wirt: Maus	Berechnete Masse: 30 kDa	
Isotyp: IgG1	Beobachtete Masse: 30 kDa	
Immunogen Katalognummer: AG32731		

Anwendungen

Geprüfte Anwendungen:
WB, ELISA

Getestete Reaktivität:
Human, Maus, Ratte

Positivkontrollen:

WB : Hausschwein-Hirngewebe, Jurkat-Zellen, K-562-Zellen, Kaninchenhirngewebe, LNCaP-Zellen, Maushirngewebe, Rattenhirngewebe

Hintergrundinformationen

TOLLIP (Toll interacting protein), is an inhibitory adaptor protein. Tollip is a negative regulator of IL-1R and Toll-like receptors signalling pathway, involved in inflammatory immune response and inhibiting IRAK1 phosphorylation and kinase activity (PMID:11441107, PMID:10854325, PMID:11751856). The TOLLIP-dependent selective autophagy pathway plays a role in clearance of cytotoxic polyQ proteins aggregates (PubMed:25042851).

Lagerung

Lagerungsbedingungen:
Bei -20°C lagern. Nach dem Versand ein Jahr lang stabil

Lagerungspuffer:
PBS mit 0.02% Natriumazid und 50% Glycerin pH 7.3.

Aliquotieren ist nicht notwendig bei -20°C Lagerung

*** 20ul-Größen enthalten 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Ausgewählte Validierungsdaten



Various lysates were subjected to SDS PAGE followed by western blot with 68170-1-Ig (TOLLIP antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Various lysates were subjected to SDS PAGE followed by western blot with 68170-1-Ig (TOLLIP antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated Beta Actin Monoclonal antibody (HRP-66009) as loading control.