

## Allgemeine Informationen

Katalog-Nr.:	GenBank-Zugangsnummer:	Reinigungsmethode:
68250-1-Ig	BC093730	Protein-A-Reinigung
Größe:	GenID (NCBI):	CloneNo.:
150ul, Konzentration: 1000 µg/ml von 5420	5420	4A3C7
Nanodrop;	Vollständiger Name:	Empfohlene Verdünnungen:
Wirt:	podocalyxin-like	IHC 1:500-1:2000
Maus	Berechneté Masse:	IF 1:200-1:800
Isotyp:	526 aa, 55 kDa	
IgG2a	Beobachteté Masse:	
Immunogen Katalognummer:	60-70 kDa	
AG12859		

## Anwendungen

Geprüfte Anwendungen:	Positivkontrollen:
FC, IF, IHC, ELISA	IHC : humanes Nierengewebe,
Getestete Reaktivität:	IF : humanes Nierengewebe,

**Hinweis-IHC: Antigendemaskierung mit TE-Puffer pH 9,0 empfohlen. (\*) Wahlweise kann die Antigendemaskierung auch mit Citratpuffer pH 6,0 erfolgen.**

## Hintergrundinformationen


Podocalyxin, also known as podocalyxin-like protein 1 (PODXL or PCLP1), is a transmembrane glycoprotein belonging to the CD34 family of sialomucins. Podocalyxin was originally identified as the major sialoprotein on podocytes of the kidney glomerulus, but was later found to be expressed on vascular endothelial cells and early hematopoietic progenitors. It involved in the regulation of both adhesion and cell morphology. In addition, podocalyxin is highly expressed in embryonic stem cells and aberrant expression of podocalyxin has been implicated in a wide range of cancers. Podocalyxin is a protein with a peptide bone of 55.5 kDa that undergoes a post-translational glycosylation, the different molecular mass of podocalyxin indicates the extent and variability of glycosylation patterns (PMID: 17092254).

## Lagerung


**Lagerungsbedingungen:**  
Bei -20°C lagern. Nach dem Versand ein Jahr lang stabil  
**Lagerungspuffer:**  
PBS mit 0.02% Natriumazid und 50% Glycerin pH 7.3.  
**Aliquotieren ist nicht notwendig bei -20°C Lagerung**

\*\*\* 20ul-Größen enthalten 0.1% BSA


## Ausgewählte Validierungsdaten




Immunohistochemical analysis of paraffin-embedded human kidney tissue slide using 68250-1-Ig (Podocalyxin antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human kidney tissue slide using 68250-1-Ig (Podocalyxin antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed human kidney tissue using Podocalyxin antibody (68250-1-Ig, Clone: 4A3C7) at dilution of 1:400 and Coralite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).



1X10<sup>6</sup> HeLa cells were intracellularly stained with 0.4 ug Anti-Human Podocalyxin (68250-1-Ig, Clone:4A3C7) and Coralite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Mouse IgG2a Isotype Control (66360-2-Ig, Clone: K11A1B2A2) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).