

Nur für Forschungszwecke

# SNX17 Monoklonaler Antikörper

Katalog-Nr.:68256-1-Ig



## Allgemeine Informationen

<b>Katalog-Nr.:</b> 68256-1-Ig	<b>GenBank-Zugangsnummer:</b> BC002524	<b>Reinigungsmethode:</b> Protein-G-Reinigung
<b>Größe:</b> 150ul , Konzentration: 500 µg/ml von Nanodrop;	<b>GeneID (NCBI):</b> 9784	<b>CloneNo.:</b> 3B8F6
<b>Wirt:</b> Maus	<b>Vollständiger Name:</b> sorting nexin 17	<b>Empfohlene Verdünnungen:</b> WB 1:5000-1:50000 IF 1:200-1:800
<b>Isotyp:</b> IgG1	<b>Berechnete Masse:</b> 53 kDa	
<b>Immunogen Katalognummer:</b> AG32624	<b>Beobachtete Masse:</b> 55 kDa	

## Anwendungen

### Geprüfte Anwendungen:

IF, WB, ELISA

### Getestete Reaktivität:

Hausschwein, Human, Ratte

### Positivkontrollen:

WB : LNCaP-Zellen, Hausschwein-Hirngewebe, HeLa-Zellen, Jurkat-Zellen, Rattenhirngewebe

IF : HepG2-Zellen, HeLa-Zellen

## Hintergrundinformationen

Sorting nexin 17 (SNX17, synonym: KIAA0064) is a member of the sorting nexin family. Members of this family contain a phox (PX) domain, which is a phosphoinositide binding domain, and are involved in intracellular trafficking. SNX17 does not contain a coiled coil region, like some family members, but contains a B41 domain. This protein interacts with the cytoplasmic domain of P-selectin, and may function in the intracellular trafficking of P-selectin.

## Lagerung

### Lagerungsbedingungen:

Bei -20°C lagern. Nach dem Versand ein Jahr lang stabil

### Lagerungspuffer:

PBS mit 0.02% Natriumazid und 50% Glycerin pH 7.3.

Aliquotieren ist nicht notwendig bei -20°C Lagerung

\*\*\* 20ul-Größen enthalten 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free  
in USA), or 1(312) 455-8498 (outside USA)


E: [proteintech@ptglab.com](mailto:proteintech@ptglab.com)  
W: [ptglab.com](http://ptglab.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Ausgewählte Validierungsdaten



Various lysates were subjected to SDS PAGE followed by western blot with 68256-1-Ig (SNX17 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using SNX17 antibody (68256-1-Ig, Clone: 3B8F6) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L), CL594-Phalloidin (red).