

Nur für Forschungszwecke

NFIC Monoklonaler Antikörper

Katalog-Nr.:68318-1-Ig



Allgemeine Informationen

Katalog-Nr.: 68318-1-Ig	GenBank-Zugangsnummer: BC012120	Reinigungsmethode: Protein-A-Reinigung
Größe: 150ul , Konzentration: 1000 µg/ml von4782 Nanodrop;	GeneID (NCBI): 4782	CloneNo.: 1A11A5
Wirt: Maus	Vollständiger Name: nuclear factor I/C (CCAAT-binding transcription factor)	Empfohlene Verdünnungen: WB 1:5000-1:50000
Isotyp: IgG2b	Berechnete Masse: 428aa,48 kDa; 508aa,57 kDa	
Immunogen Katalognummer: AG9472	Beobachtete Masse: 50-56 kDa	

Anwendungen

Geprüfte Anwendungen: WB, ELISA	Positivkontrollen: WB : HSC-T6 cells, 4T1-Zellen, HeLa-Zellen, LNCaP-Zellen, NIH/3T3-Zellen
Getestete Reaktivität: Human, Maus, Ratte	

Hintergrundinformationen

NFIC, also named as Nuclear factor 1 C-type, is a 508 amino acid protein, which contains 1 CTF/NF-I DNA-binding domain and belongs to the CTF/NF-I family. NFIC localizes in the nucleus and recognizes and binds the palindromic sequence 5'-TTGGCNNNNNGCCAA-3' present in viral and cellular promoters and in the origin of replication of adenovirus type 2. These proteins are individually capable of activating transcription and replication.

Lagerung

Lagerungsbedingungen:
Bei -20°C lagern. Nach dem Versand ein Jahr lang stabil
Lagerungspuffer:
PBS mit 0.02% Natriumazid und 50% Glycerin pH 7.3.
Aliquotieren ist nicht notwendig bei -20°C Lagerung

*** 20ul-Größen enthalten 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Ausgewählte Validierungsdaten



Various lysates were subjected to SDS PAGE followed by western blot with 68318-1-Ig (NFIC antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours.