

Nur für Forschungszwecke

NeutraKine® SCF Monoklonaler Antikörper

Katalog-Nr.: 69026-1-Ig



Allgemeine Informationen

Katalog-Nr.:

69026-1-Ig

Größe:

100ug

Wirt:

Maus

Isotyp:

IgG1

Immunogen Katalognummer:

HZ-1024

GenBank-Zugangsnummer:

GeneID (NCBI):

4254

Vollständiger Name:

KIT ligand

Reinigungsmethode:

Protein-G-Reinigung

CloneNo.:

1B3C4

Anwendungen

Geprüfte Anwendungen:

Neutralization, ELISA

Getestete Reaktivität:

Human

Hintergrundinformationen

Stem cell factor (SCF) also known as mast cell growth factor, steel factor, and kit ligand, is a hematopoietic growth factor. SCF is a critical protein with key roles in the cell such as hematopoiesis, gametogenesis and melanogenesis. SCF activates multiple signal pathways by binding with c-kit receptor. SCF and its receptor, c-kit ligand (KL), are up-regulated in particular human malignancies including gastrointestinal stromal tumor (GISTs), breast cancer, hematopoietic cell, myeloid leukaemia, and glioma. The SCF/c-KIT system also plays a relevant role in cell fate decision, and its activity is widely known to control cell proliferation, differentiation, survival and apoptosis.

This antibody is used to neutralize the bioactivity of SCF.

Lagerung

Lagerungsbedingungen:

Lyophilisierte Antikörper sind bei Lagerung zwischen (-20°C) und (-80°C) 1 Jahr ab Empfangsdatum stabil. Bei Rekonstitution empfehlen wir, die Lösung kurzfristig bei (4°C) oder langfristig bei (-20°C) bis (-80°C). Rekonstituierte Produkte sollten nicht wiederholt eingefroren und wieder aufgetaut werden.

Lagerungspuffer:

Sterile PBS.

Aliquotieren ist nicht notwendig bei -20°C Lagerung

*** 20ul-Größen enthalten 0.1% BSA

For technical support and original validation data for this product please contact:


T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com

W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Ausgewählte Validierungsdaten



Recombinant human SCF (Cat.NO. HZ-1024) stimulates proliferation of TF-1 cell line (human erythroleukemic cell line) in a dose-dependent manner (blue curve, refer to bottom X-left Y). The activity of human SCF (20 ng/mL HZ-1024) is neutralized by mouse anti-human SCF monoclonal antibody 69026-1-Ig at serial dose (red curve, refer to top X-right Y). The ND50 is typically 0.5-2 µg/mL.