

# NeutraControl Noggin Monoklonaler Antikörper

Katalog-Nr.: 69510-1-Ig

## Allgemeine Informationen

Katalog-Nr.:	GenBank-Zugangsnummer:	Reinigungsmethode:
69510-1-Ig	GenID (NCBI):	Protein-A-Reinigung
Größe:	9241	CloneNo.:
100ug, Konzentration: 1000 µg/ml von Nanodrop;	Vollständiger Name:	3B10F3
Wirt:	noggin	
Maus		
Isotyp:		
IgG2a		
Immunogen Katalognummer:		
HZ-1118		

## Anwendungen

Geprüfte Anwendungen:  
Non-Neutralization, ELISA

Getestete Reaktivität:  
Human

## Hintergrundinformationen

Noggin is an extracellular polypeptide acting as an antagonist of bone morphogenetic proteins (BMPs) regulating embryonal development. Noggin inhibits activity of BMP-2, -4, -7, -13, and -14. Noggin is present extracellularly in the matrix or retained at the cell surface via interaction with heparin sulfate proteoglycans. In early development stages, Noggin is produced by the Spemann organizer, allowing dorsal-ventral patterning of BMPs (PMID: 8752214). Subsequently, Noggin is expressed by the notochord regulating BMP-4 signaling in neurogenesis. Additionally, Noggin is present during development in the dermal papilla, connective tissue of the hair follicle, lens, retina, and periocular mesenchyme, as well as in the mesoderm lineage regulating development of the bone, cartilage, and muscles.


This antibody is a neutralizing control antibody for Noggin, the immunogen is the same as Neutrakine 69010-1-Ig but could not neutralize human Noggin.

## Lagerung

**Lagerungsbedingungen:**  
Lyophilisierte Antikörper sind bei Lagerung zwischen (-20°C) und (-80°C) 1 Jahr ab Empfangsdatum stabil. Bei Rekonstitution empfehlen wir, die Lösung kurzfristig bei (4°C) oder langfristig bei (-20°C) bis (-80°C). Rekonstituierte Produkte sollten nicht wiederholt eingefroren und wieder aufgetaut werden.  
**Lagerungspuffer:**  
Sterile PBS.  
**Aliquotieren ist nicht notwendig bei -20°C Lagerung**

\*\*\* 20ul-Größen enthalten 0.1% BSA

## Ausgewählte Validierungsdaten



Recombinant human Noggin (Cat.NO. HZ-1118) inhibits BMP-4 induced alkaline phosphate production (using pNPP as chromogenic substrate for detection) by ATDC5 in a dose dependent manner (blue curve, refer to bottom X-Left Y). The activity of human Noggin (40 ng/mL HZ-1118) is neutralized by mouse anti-human Noggin monoclonal antibody 69010-1-Ig at serial dose (red curve, refer to top X-right Y). The ND<sub>50</sub> is typically 5-20 µg/mL. The



Indirect ELISA was carried out by coating recombinant Human Noggin (Cat.NO. HZ-1118) at 70 ng/well followed by blocking and adding serial diluted Noggin antibody 69010-1-Ig and 69510-1-Ig respectively. Signal was developed with TMB and stopped by H<sub>2</sub>SO<sub>4</sub>. Signal strength was measured by absorbance at 450 nm.