

Nur für Forschungszwecke

# NeutraControl VEGF165 Monoklonaler Antikörper



Katalog-Nr.: 69525-1-Ig

## Allgemeine Informationen

Katalog-Nr.: 69525-1-Ig	GenBank-Zugangsnummer: GeneID (NCBI): 7422	Reinigungsmethode: Protein-G-Reinigung
Größe: 100ug	Vollständiger Name: vascular endothelial growth factor A	CloneNo.: 7E9F4
Wirt: Maus		
Isotyp: IgG1		
Immunogen Katalognummer: HZ-1038		

## Anwendungen

**Geprüfte Anwendungen:**  
Non-Neutralization, ELISA

**Getestete Reaktivität:**  
Human

## Hintergrundinformationen

VEGFA, also named as VEGF or VPF, belongs to the PDGF/VEGF growth factor family. It is a growth factor active in angiogenesis, vasculogenesis and endothelial cell growth. VEGFA induces endothelial cell proliferation, promotes cell migration, inhibits apoptosis and induces permeabilization of blood vessels. It binds to the FLT1/VEGFR1 and KDR/VEGFR2 receptors, heparan sulfate and heparin. Defects in VEGFA are associated with microvascular complications of diabetes type 1 (MVCD1). VEGFA has 17 isoforms with MW from 16 to 45 kDa. Some isoforms have homodimer forms (e.g.; VEGFA189 38 kDa or VEGFA110 34 kDa). VEGF-A exists in at least seven homodimeric isoforms. The monomers consist of 121, 145, 148, 165, 183, 189, or 206 amino acids (PMID:15602010).

This antibody is a neutralizing control antibody for VEGF165, the immunogen is the same as Neutrakine 69025-1-Ig but could not neutralize human VEGF165.

## Lagerung

**Lagerungsbedingungen:**  
Lyophilisierte Antikörper sind bei Lagerung zwischen (-20°C) und (-80°C) 1 Jahr ab Empfangsdatum stabil. Bei Rekonstitution empfehlen wir, die Lösung kurzfristig bei (4°C) oder langfristig bei (-20°C) bis (-80°C). Rekonstituierte Produkte sollten nicht wiederholt eingefroren und wieder aufgetaut werden.

**Lagerungspuffer:**  
Sterile PBS.

Aliquotieren ist nicht notwendig bei -20°C Lagerung

\*\*\* 20ul-Größen enthalten 0.1% BSA



For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: [proteintech@ptglab.com](mailto:proteintech@ptglab.com)  
W: [ptglab.com](http://ptglab.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Ausgewählte Validierungsdaten



Recombinant human VEGF165 (Cat.NO. HZ-1038) stimulates proliferation of HUVEC (human umbilical vein endothelial) cell line in a dose-dependent manner (blue curve, refer to bottom X-left Y axis). The activity of human VEGF165 (10 ng/mL HZ-1038) is neutralized by mouse anti-human VEGF165/121 monoclonal antibody 69025-1-Ig at serial dose (red curve, refer to top X-right Y axis). The ND50 is typically 0.5-2 µg/mL. The NeutraControl mouse anti-human VEGF165



Indirect ELISA was carried out by coating recombinant Human VEGF165 (Cat.NO. HZ-1038) and VEGF121 (Cat.NO. HZ-1204) respectively at 70 ng/well followed by blocking and adding serial diluted VEGF165/121 antibody 69025-1-Ig and VEGF165 antibody 69525-1-Ig respectively. HRP-goat anti-mouse was used for detection. Signal was developed with TMB and stopped by H<sub>2</sub>SO<sub>4</sub>. Signal strength was measured by absorbance at 450 nm. The result suggests that

