

Allgemeine Informationen

Katalog-Nr.: 80001-1-RR	GenBank-Zugangsnummer: BC001487	Reinigungsmethode: Protein-A-Reinigung
Größe: 100ul, Konzentration: 250 µg/ml von Nanodrop;	GeneID (NCBI): 23435	CloneNo.: 11N20
Wirt: Kaninchen	Vollständiger Name: TAR DNA binding protein	Empfohlene Verdünnungen: WB 1:5000-1:50000 IHC 1:500-1:4000 IF 1:300-1:1200
Isotyp: IgG	Berechnete Masse: 43 kDa	
	Beobachtete Masse: 43 kDa	

Anwendungen

Geprüfte Anwendungen:

FC, IF, IHC, WB, ELISA

Getestete Reaktivität:

Human, Maus, Ratte

Hinweis-IHC: Antigendemaskierung mit TE-Puffer pH 9,0 empfohlen. (*) Wahlweise kann die Antigendemaskierung auch mit Citratpuffer pH 6,0 erfolgen.

Positivkontrollen:

WB: HeLa-Zellen, C6-Zellen, K-562-Zellen, Neuro-2a-Zellen

IHC: humanes Gliomgewebe, Maushirngewebe, Rattenhirngewebe

IF: Rattenhirngewebe, HAP1 cells

Hintergrundinformationen

The TARDBP gene encodes the TDP-43 protein, initially found to repress HIV-1 transcription by binding TAR DNA. TDP-43 has since been shown to bind RNA as well as DNA, and have multiple functions in transcriptional repression, translational regulation and pre-mRNA splicing. For instance, it is reported to regulate alternate splicing of the CTFE gene. In 2006 Neumann et al. found that hyperphosphorylated, ubiquitinated and/or cleaved forms of TDP-43, collectively known as pathological TDP-43, play a major role in the disease mechanisms of ubiquitin-positive, tau- and alpha-synuclein-negative frontotemporal dementia (FTLD-U) and in amyotrophic lateral sclerosis (ALS). Proteintech's 80001-1-RR is a rabbit recombinant TDP-43 antibody recognizing N-terminal TDP-43. It recognizes the intact 43 kDa protein as well as all posttranslationally modified and truncated forms in multiple applications. Various forms of TDP-43 exist, including 18-35 kDa of cleaved C-terminal fragments, 45-50 kDa phospho-protein, 55 kDa glycosylated form, 75 kDa hyperphosphorylated form, and 90-300 kDa cross-linked form. (PMID: 17023659, 19823856, 21666678, 22193176) Recently TDP-43 has been reported to be overexpressed in triple negative breast cancer (TNBC) and it may be a potential target for TNBC diagnosis and drug design. (PMID: 29581274).

80001-1-RR can be used in WB, IHC, IHF. For the ICC (IF with cells) experiment, the results will be better with ethanol fixed cell and dilution 1:50 or 1:100. Another antibody with CatNo. 80002-1-RR is recommended for IF or ICC experiment.

Lagerung

Lagerungsbedingungen:

Bei -20°C lagern. Nach dem Versand ein Jahr lang stabil

Lagerungspuffer:

PBS mit 0.02% Natriumazid und 50% Glycerin pH 7.3.

Aliquotieren ist nicht notwendig bei -20°C Lagerung

*** 20ul-Größen enthalten 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Ausgewählte Validierungsdaten




Various cell lysates were subjected to SDS PAGE followed by western blot with 80001-1-RR (TDP43 antibody) at dilution of 1:35000 incubated at room temperature for 1.5 hours.




HAP1 (WT and TARDBP KO) lysates prepared with NP-40 buffer, 50 µg protein loaded. 80001-1-RR incubated at 1:1000 at 4°C overnight in 5% milk in TBST. Ponceau stained transfers shown on right. Data provided by YCharOS, an open science company with a mission to validate commercial antibodies to improve scientific reproducibility and transparency.




WB result of TDP-43 antibody (80001-1-RR; 1:30000; incubated at room temperature for 1.5 hours) with sh-Control and sh-TDP-43 transfected HeLa cells.




Immunofluorescent analysis of (4% PFA) fixed rat brain tissue using 80001-1-RR (TDP-43 antibody) at dilution of 1:150 and CoralLite488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).




HAP1 WT cells (yellow outline) and TARDBP KO cells (red outline) labelled with a green or a far red fluorescence dye, respectively. Cells fixed with 4% PFA and stained with 80001-1-RR at 1:400 plus DAPI. Bars = 10 µm. Data provided by YCharOS, an open science company with a mission to validate commercial antibodies to improve scientific reproducibility and transparency.




IHC results of TDP43 rabbit recombinant antibody (80001-1-RR, 1:20000) with the spinal cord slides from ALS patients. IHC experiment was done with Ventana automatic staining system and Optiview DAB detection kit with heat-induced epitope retrieval (boiling for 32 min in Tris-EDTA based solution CC1 buffer, Ventana). Fig from the lab of Dr. Neumann.





IHC results of TDP43 rabbit recombinant antibody (80001-1-RR, 1:20000) with the frontal cortex from FTLD-TDP type B patients. IHC experiment was done with Ventana automatic staining system and Optiview DAB detection kit with heat-induced epitope retrieval (boiling for 32 min in Tris-EDTA based solution CC1 buffer, Ventana). Fig from the lab of Dr. Neumann.



Immunohistochemical analysis of paraffin-embedded human gliomas tissue slide using 80001-1-RR (TDP-43 antibody) at dilution of 1:1500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human gliomas tissue slide using 80001-1-RR (TDP-43 antibody) at dilution of 1:1500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded rat brain tissue slide using 80001-1-RR (TDP-43 antibody) at dilution of 1:2000 (under 4x lens).

1X10⁶ HeLa cells were intracellularly stained with 0.20ug Anti-Human TDP-43 (80001-1-RR, Clone:11N20) (red) or 0.20 ug control antibody (blue) and CoraLite488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) with dilution 1:1000. Fixed with 90% MeOH.