

# C1QBP Rekombinanter Antikörper

Katalog-Nr.:**80490-1-RR**

## Allgemeine Informationen

Katalog-Nr.:	GenBank-Zugangsnummer:
80490-1-RR	BC013731
Größe:	GenID (NCBI):
100ul , Konzentration: 1000 µg/ml von 708	
Nanodrop;	Vollständiger Name:
	complement component 1, q
Wirz:	subcomponent binding protein
Kaninchen	
Isotyp:	Berechneté Masse:
IgG	282 aa, 31 kDa
Immunogen Katalognummer:	Beobachteté Masse:
AG19773	32 kDa

Reinigungsmethode:  
Protein-A-Reinigung

CloneNo.:

3K20

Empfohlene Verdünnungen:  
WB 1:2000-1:10000  
IHC 1:50-1:500  
IF 1:250-1:1000

## Anwendungen

Geprüfte Anwendungen:  
FC, IF, IHC, WB, ELISA

Getestete Reaktivität:  
Human, Maus, Ratte

**Hinweis-IHC:** Antigendemaskierung mit **TE-Puffer pH 9,0 empfohlen.** (\*) Wahlweise kann die Antigendemaskierung auch mit **Citratpuffer pH 6,0 erfolgen.**

Positivkontrollen:

WB : HeLa-Zellen, 4T1-Zellen, HEK-293-Zellen, HepG2-Zellen, Jurkat-Zellen, K-562-Zellen, NIH/3T3-Zellen, PC-12-Zellen

IHC : humanes Tonsillitisgewebe,  
IF : HeLa-Zellen,

## Hintergrundinformationen


C1QBP, also named as gC1q receptor (gC1qR), p32, p33, and hyaluronan-binding protein 1 (HABP1), is a protein initially copurified with splicing factor SF2 (PMID: 1830244). The protein is synthesized as a pro-protein of 282 amino acids (aa) that is post-translationally processed by removal of the initial 73 aa to a mature protein of 209 aa (PMID: 8262387). C1QBP is an evolutionary conserved and ubiquitously expressed multifunctional protein and has been reported to be a predominantly mitochondrial matrix protein involved in inflammation and infection processes, mitochondrial ribosome biogenesis, regulation of apoptosis and nuclear transcription, and pre-mRNA splicing (PMID: 28942965).

## Lagerung





Lagerungsbedingungen:  
Bei -20°C lagern. Nach dem Versand ein Jahr lang stabil  
Lagerungspuffer:  
PBS mit 0,02% Natriumazid und 50% Glycerin pH 7,3.  
Aliquotieren ist nicht notwendig bei -20°C Lagerung

\*\*\* 20ul-Größen enthalten 0,1% BSA

## Ausgewählte Validierungsdaten



Various lysates were subjected to SDS PAGE followed by western blot with 80490-1-RR (C1QBP antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



1X10<sup>6</sup> HeLa cells were intracellularly stained with 0.5 ug Anti-Human C1QBP (80490-1-RR, Clone:3K20) (red) labeled with FlexAble CoraLite® Plus 555 Antibody Labeling Kit for Rabbit IgG (KFA002), or 0.5 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).