## Nur für Forschungszwecke

## CD27 Monoklonaler Antikörper

Katalog-Nr.:AB-65132



Allgemeine Informationen

Katalog-Nr.: GenBank-Zugangsnummer: AB-65132 BC012160

GeneID (NCBI):

100tests , 5 µl/test 93

Wirt: Vollständiger Name:
Maus CD27 molecule
Isotyp: Berechneté Masse:

IgG1, kappa 29 kDa

Reinigungsmethode: Affinitätsreinigung

CloneNo.: 0323

Anregungs-/Emissionsmaxima-

Wellenlängen: 404 nm / 458 nm

Anwendungen

Geprüfte Anwendungen:

FC

Größe:

Getestete Reaktivität:

Human

Hintergrundinformationen

CD27 (also known as TNFRSF7) is a type I glycoprotein expressed on some B cells and the majority of T cells, and is a member of the tumor necrosis factor (TNF) receptor family. CD27 is required for generation and long-term maintenance of T cell immunity (PMID: 11062504). It is a receptor for CD70 (CD27L). Ligation of CD27 by CD70 induces strong ubiquitination of TRAF and the activation of both canonical and non-canonical NF-kappaB pathways, as well as the JNK pathway (PMID: 19426224). CD27 may also play a role in apoptosis through association with SIVA1.

Lagerung


Lagerungsbedingungen:

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.


Lagerungspuffer:

PBS with 0.09% sodium azide and 0.5% BSA.

## Ausgewählte Validierungsdaten



1X10^6 human PBMCs were surface stained with 5 ul Atlantic Blue™ Anti-Human CD27 (AB-65132, Clone:0323) (red) or Mouse IgG1 Isotype Control. Cells were not fixed. Lymphocytes were gated.



1X10^6 human PBMCs were surface stained with 5 ul Atlantic Blue™ Anti-Human CD27 (AB-65132, Clone:O323) or Mouse IgG1 Isotype Control. Cells were not fixed. Lymphocytes were gated.