Nur für Forschungszwecke

## CD73 Monoklonaler Antikörper

Katalog-Nr.:AB-65162





Allgemeine Informationen	Katalog-Nr.: AB-65162	GenBank-Zugangsnummer: BC015940	Reinigungsmethode: Protein-G-Reinigung
	Größe: 100tests , 5 µl/test	GenelD (NCBI): 4907	CloneNo.: AD2
	Wirt: Maus Isotyp: IgG1, kappa	Vollständiger Name: 5'-nucleotidase, ecto (CD73) Berechneté Masse: 29 kDa, 63 kDa	Anregungs-/Emissionsmaxima- Wellenlängen: 404 nm / 458 nm
Getestete Reaktivität: Human			
Hintergrundinformationen	CD73, also known as ecto-5'-nucleotidase (5'-NT), is a 70-kDa, glycosyl-phosphatidylinositol-linked membrane- bound glycoprotein found in most tissues (PMID: 18404475; 20179192). CD73 is an ectoenzyme that catalyzes the dephosphorylation of AMP and other nucleoside monophosphates (PMID: 9553767). In the human immune system, CD73 is expressed on subsets of T and B cells, on germinal center follicular dendritic cells, and on thymic medullary reticular fibroblasts and epithelial cells (PMID: 2137649; 9553767). CD73 is highly expressed in many human solid tumors and is closely involved in cancer progression (PMID: 20179192).		
Lagerung	Lagerungsbedingungen: Store at 2-8°C. Avoid exposur Lagerungspuffer: PBS with 0.09% sodium azide	e to light. Stable for one year after shipme a and 0.5% BSA.	ent.

For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll freeE: proteintech@ptglab.comin USA), or 1(312) 455-8498 (outside USA)W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Ausgewählte Validierungsdaten





1X10^6 human PBMCs were surface co-stained with FITC Anti-Human CD3 and 5 ul Atlantic Blue™ Anti-Human CD73 (AB-65162, Clone:AD2) or Mouse IgG1 Isotype Control. Cells were not fixed. Lymphocytes were gated.

1X10<sup>^</sup>6 human PBMCs were surface stained with 5 ul Atlantic Blue<sup>™</sup> Anti-Human CD73 (AB-65162, Clone:AD2) or Mouse IgG1 Isotype Control. Cells were not fixed. Lymphocytes were gated.