Nur für Forschungszwecke

CD8 Monoklonaler Antikörper

Katalog-Nr.:AB-65204



Allgemeine Informationen

Katalog-Nr.:

AB-65204 Größe:

100 tests , 5 μ l/test

Wirt: Maus Isotyp:

lgG2a

GenBank-Zugangsnummer:

BC025715 GeneID (NCBI):

Vollständiger Name: CD8a molecule Berechneté Masse:

235 aa, 26 kDa

Reinigungsmethode:

Protein-A-Reinigung

CloneNo.: UCHT4

Anregungs-/Emissionsmaxima-

Wellenlängen: 404 nm / 458 nm

Anwendungen

Geprüfte Anwendungen:

FC

Getestete Reaktivität:

Human

Hintergrundinformationen

CD8 is a transmembrane glycoprotein composed of two disulfide-linked chains. It can be present as a homodimer of CD8a or as a heterodimer of CD8a and CD8β (PMID: 3264320; 8253791). CD8 is found on most thymocytes. The majority of class I-restricted T cells express mostly the CD8αβ heterodimer while CD8αα homodimers alone have been found on some gut intraepithelial T cells, on some T cell receptor (TCR) γδ T cells and on NK cells (PMID: 2111591; 1831127; 8420975). CD8 acts as a co-receptor that binds to MHC class-I and participates in cytotoxic T cell activation (PMID: 8499079). During T cell development, CD8 is required for positive selection of CD4-/CD8+T cells (PMID: 1968084).

Lagerung


Lagerungsbedingungen:

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.


Lagerungspuffer:

PBS with 0.09% sodium azide and 0.5% BSA.

Ausgewählte Validierungsdaten



1X10^6 human PBMCs were surface co-stained with APC Anti-Human CD3 and 5 ul Atlantic Blue™ Anti-Human CD8 (AB-65204, Clone: UCHT4) or Mouse IgG1 Isotype Control. Cells were not fixed. Lymphocytes were gated.



1X10^6 human PBMCs were surface stained with 5 ul Atlantic Blue™ Anti-Human CD8 (AB-65204, Clone: UCHT4) or Mouse IgG1 isotype control. Cells were not fixed. Lymphocytes were gated.