

Nur für Forschungszwecke

# CD107b / LAMP2 Monoklonaler Antikörper

Katalog-Nr.:[APC-65052](#)



## Allgemeine Informationen

Katalog-Nr.:	GenBank-Zugangsnummer:	Reinigungsmethode:
APC-65052	BC138718	Affinitätsreinigung
Größe:	GenID (NCBI):	CloneNo.:
100ug , 0.1 mg/ml	16784	ABL-93
Wirt:	Vollständiger Name:	Empfohlene Verdünnungen:
Ratte	lysosomal-associated membrane protein 2	IF 1:50-1:500
Isotyp:		Anregungs-/Emissionsmaxima-Wellenlängen:
IgG2a		650 nm / 660 nm

## Anwendungen

Geprüfte Anwendungen:	Positivkontrollen:
FC (Intra), IF	IF : NIH/3T3-Zellen,
Getestete Reaktivität:	
Maus	

## Hintergrundinformationen

LAMP2 (CD107b) is a Lysosomal membrane glycoprotein. LAMP2 is extensively glycosylated with asparagine-linked oligosaccharides which protect it from intracellular proteolysis (PMID: 10521503). Although LAMP-2 is localized primarily in the endosome-lysosomal membrane of cells, it is also found on the plasma membrane under certain circumstances, e.g., after platelet activation, during granulocytic differentiation and activation, and in some tumor cells (PMID: 12221139). LAMP is involved in lysosomal stability and autophagy (PMID: 12221139). This glycoprotein provides selectins with carbohydrate ligands. LAMP2 may play a role in tumor cell metastasis (PMID 9426697).

## Lagerung


Lagerungsbedingungen:  
Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.  
Lagerungspuffer:  
PBS mit 0,1% Natriumazid und einem Stabilisator.

For technical support and original validation data for this product please contact:  
T: 1(888) 4PTGLAB (1-888-478-4522) (toll free  
in USA), or 1(312) 455-8498 (outside USA)


E: [proteintech@ptglab.com](mailto:proteintech@ptglab.com)  
W: [ptglab.com](http://ptglab.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Ausgewählte Validierungsdaten



Immunofluorescent analysis of (4% PFA) fixed NIH/3T3 cells using APC-65052 (CD107b antibody) at dilution of 1:100.



$1 \times 10^6$  NIH/3T3 cells were intracellularly stained with 10  $\mu$ l APC Anti-Mouse CD107b (APC-65052, Clone: ABL-93) (red), or 10  $\mu$ l Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).