

Nur für Forschungszwecke

CD11c Monoklonaler Antikörper

Katalog-Nr.: APC-65086



Allgemeine Informationen

| | | |
|---------------------------------------|---|--|
| Katalog-Nr.: APC-65086 | GenBank-Zugangsnummer: BC038237 | Reinigungsmethode: Affinitätsreinigung |
| Größe: 100tests , 5 µl/test | GeneID (NCBI): 3687 | CloneNo.: 3,9 |
| Wirt: Maus | Vollständiger Name: integrin, alpha X (complement component 3 receptor 4 subunit) | Anregungs-/Emissionsmaxima-Wellenlängen: 650 nm / 660 nm |
| Isotyp: IgG1, kappa | Berechnete Masse: 1169 aa, 129 kDa | |

Anwendungen

Geprüfte Anwendungen:
FC

Getestete Reaktivität:
Human

Hintergrundinformationen

Integrins are cell adhesion receptors that are heterodimers composed of non-covalently associated α and β subunits (PMID: 9779984). CD11c, also known as integrin αX , is a type I transmembrane glycoprotein present on a variety of cells, including monocytes/macrophages, granulocytes, a subset of B cells, NK cells and dendritic cells (PMID: 2897326; 1680915; 1694698; 17389580). As a result of its high level of expression on most dendritic cells, CD11c is typically considered to be a marker of conventional dendritic cells (PMID: 27119555). CD11c forms an α/β heterodimer with CD18 (integrin $\beta 2$). CD11c/CD18 acts a receptor for fibrinogen and is important in monocyte adhesion and chemotaxis (PMID: 1671533).

Lagerung

Lagerungsbedingungen:
Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.
Lagerungspuffer:
Phosphatbasierter Puffer mit 0,09% Natriumazid und 0,1% Gelatine, pH 7,2.


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Ausgewählte Validierungsdaten



100 ul human peripheral blood were surface stained with 5.00 ul APC Anti-Human CD11c (APC-65086, Clone:3.9) and then treated with red blood cell lysis buffer. Cells were not fixed.



100 ul human peripheral blood were surface stained with 0.25 ug APC-mouse IgG1 isotype control and then treated with red blood cell lysis buffer. Cells were not fixed.