

Nur für Forschungszwecke

CD19 Monoklonaler Antikörper

Katalog-Nr.: APC-65145



Allgemeine Informationen

Katalog-Nr.: APC-65145	GenBank-Zugangsnummer: BC006338	Reinigungsmethode: Affinitätsreinigung
Größe: 100tests, 5 µl(0.25µg)/test	GeneID (NCBI): 930	CloneNo.: SJ25C1
Wirt: Maus	Vollständiger Name: CD19 molecule	Anregungs-/Emissionsmaxima-Wellenlängen: 650 nm / 660 nm
Isotyp: IgG1	Berechnete Masse: 556 aa, 61 kDa	

Anwendungen

Geprüfte Anwendungen:
FC

Getestete Reaktivität:
Human

Hintergrundinformationen

CD19 is a 95 kDa type I transmembrane glycoprotein belonging to the immunoglobulin superfamily (PMID: 2472450). It is expressed by B cells and follicular dendritic cells. CD19 is up-regulated at the step of B-lineage commitment during the differentiation of the hematopoietic stem cell, it remains on during subsequent stages of differentiation until finally down-regulated during terminal differentiation into plasma cells (PMID: 8528044). CD19 is involved in B cell development, activation and differentiation. It is the dominant component for the signaling complex on B cells that includes CD21 (CR2), CD81 (TAPA-1) and CD225 and acts as a critical co-receptor for BCR signal transduction (PMID: 23210908).

Lagerung

Lagerungsbedingungen:
Bei 2-8°C lagern. Vor Licht schützen.

Lagerungspuffer:
Phosphatbasierter Puffer mit 0,09% Natriumazid und 0,1% Gelatine, pH 7,2.


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Ausgewählte Validierungsdaten



100 ul human peripheral blood were surface stained with FITC Anti-Human CD3 (FITC-65151, Clone: UCHT1) and 5.00 ul APC Anti-Human CD19 (APC-65145, Clone: SJ25C1). Lymphocytes were gated for analysis. Cells were not fixed.



100 ul human peripheral blood were surface stained with FITC Anti-Human CD3 (FITC-65151, Clone: UCHT1) and APC Mouse IgG1 isotype control antibody. Lymphocytes were gated for analysis. Cells were not fixed.