

# CD73 Monoklonaler Antikörper

Katalog-Nr.:[APC-65162](#)

## Allgemeine Informationen

<b>Katalog-Nr.:</b>	<b>GenBank-Zugangsnummer:</b>	<b>Reinigungsmethode:</b>
APC-65162	BC015940	Der gereinigte Antikörper ist mit Allophycocyanin (APC) unter optimalen Bedingungen konjugiert.
<b>Größe:</b>	<b>GenID (NCBI):</b>	Das Konjugat wird mit Größenausschluss-Chromatographie gereinigt.
100tests, 10 µl/test	4907	
<b>Wirt:</b>	<b>Vollständiger Name:</b>	
Maus	5'-nucleotidase, ecto (CD73)	
<b>Isotyp:</b>	<b>Berechnete Masse:</b>	
Mouse IgG1 kappa	29 kDa, 63 kDa	
		<b>CloneNo.:</b>
		AD2
		<b>Anregungs-/Emissionsmaxima-Wellenlängen:</b>
		650 nm / 660 nm

## Anwendungen

Geprüfte Anwendungen:

FC

Getestete Reaktivität:

Human

## Hintergrundinformationen

CD73, also known as ecto-5'-nucleotidase (5'-NT), is a 70-kDa, glycosyl-phosphatidylinositol-linked membrane-bound glycoprotein found in most tissues (PMID: 18404475; 20179192). CD73 is an ectoenzyme that catalyzes the dephosphorylation of AMP and other nucleoside monophosphates (PMID: 9553767). In the human immune system, CD73 is expressed on subsets of T and B cells, on germinal center follicular dendritic cells, and on thymic medullary reticular fibroblasts and epithelial cells (PMID: 2137649; 9553767). CD73 is highly expressed in many human solid tumors and is closely involved in cancer progression (PMID: 20179192).

## Lagerung


Lagerungsbedingungen:

Bei 2-8°C lagern. Vor Licht schützen.

Lagerungspuffer:

PBS mit 0,1% Natriumazid.

## Ausgewählte Validierungsdaten



1X10<sup>6</sup> human peripheral blood lymphocytes were surface stained with PE-Anti-Human CD3 (PE-65148, clone SK7) and 10.00 ul/test APC-Anti-Human CD73 (APC-65162, Clone:AD2). Cells were not fixed.

1X10<sup>6</sup> human peripheral blood lymphocytes were surface stained with PE-Anti-Human CD3 (PE-65148, clone SK7) and APC-mouse IgG1 isotype control antibody. Cells were not fixed.