

Nur für Forschungszwecke

GORASP2 Polyklonaler Antikörper

Katalog-Nr.:CL488-10598

Vorgestelltes Produkt



Allgemeine Informationen

Katalog-Nr.: CL488-10598	GenBank-Zugangsnummer: BC007770	Reinigungsmethode: Antigen-Affinitätsreinigung
Größe: 100ul , Konzentration: 1000 µg/ml von26003 Nanodrop;	GeneID (NCBI): Vollständiger Name: golgi reassembly stacking protein 2, 55kDa	Empfohlene Verdünnungen: IF 1:400-1:1600
Wirt: Kaninchen	Berechnete Masse: 55 kDa	Anregungs-/Emissionsmaxima-Wellenlängen: 493 nm / 522 nm
Isotyp: IgG	Beobachtete Masse: 59 kDa	
Immunogen Katalognummer: AG0950		

Anwendungen

Geprüfte Anwendungen: FC (Intra), IF	Positivkontrollen: IF : HepG2-Zellen,
Getestete Reaktivität: Human, Maus, Ratte	

Hintergrundinformationen

GORASP2 (Golgi reassembly-stacking protein 2, or GRS2) has multiple phosphorylation sites and can be phosphorylated by MKK/ERK pathway during mitosis. GORASP2 plays a role in the assembly and membrane stacking of the Golgi cisternae, and in the process by which Golgi stacks reform after mitotic breakdown. GORASP2 is supposed to regulate the intracellular transport and presentation of a defined set of transmembrane proteins, such as transmembrane TGFA.

Lagerung

Lagerungsbedingungen:
Bei -20°C lagern. Vor Licht schützen. Nach dem Versand ein Jahr stabil.
Lagerungspuffer:
BS mit 50% Glycerin, 0,05% Proclin300, 0,5% BSA, pH 7,3.
Aliquotieren ist nicht notwendig bei -20°C Lagerung

*** 20ul-Größen enthalten 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Ausgewählte Validierungsdaten



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using Coralite® Plus 488 GORASP2 antibody (CL488-10598) at dilution of 1:800, CL594-Phalloidin (red).



1×10^6 HepG2 cells were intracellularly stained with 0.4 μ g Coralite® Plus 488 Anti-Human GORASP2 (CL488-10598) (red), or 0.4 μ g Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).