

# CD25 Monoklonaler Antikörper

Katalog-Nr.: CL488-65096

## Allgemeine Informationen

Katalog-Nr.:	CL488-65096	GenBank-Zugangsnummer:	BC156627	Reinigungsmethode:
Größe:	100tests, 5 µl/test	GenID (NCBI):	3559	Affinitätsreinigung
Wirt:	Maus	Vollständiger Name:	interleukin 2 receptor, alpha	CloneNo.:
Isotyp:	IgG1	Berechneté Masse:	272 aa, 31 kDa	Anregungs-/Emissionsmaxima-Wellenlängen: 493 nm / 522 nm

## Anwendungen

Geprüfte Anwendungen:  
FC

Getestete Reaktivität:  
Human


## Hintergrundinformationen

Proliferation of T lymphocytes is triggered by the interaction of IL-2 with its specific receptor following T lymphocyte activation. The receptor for IL-2 has three forms, generated by different combinations of three different proteins, the alpha chain (IL-2R alpha), the beta chain (IL-2R beta), and the gamma chain (IL-2R gamma) (PMID: 8476561). IL-2R alpha (also known as CD25) is a type I transmembrane protein present on activated T cells, activated B cells, some thymocytes, myeloid precursors, and oligodendrocytes. CD25 is also found on natural CD4+Foxp3+ Treg cells and a subset of human CD4+ memory T cells (PMID: 22585674).


## Lagerung

Lagerungsbedingungen:  
Bei 2-8°C lagern. Vor Licht schützen.  
Lagerungspuffer:  
PBS mit 0,1% Natriumazid und 0,5% BSA, pH 7,3.

## Ausgewählte Validierungsdaten



1X10<sup>6</sup> human peripheral blood mononuclear cells (PBMCs) were surface stained with APC Anti-Human CD4 (APC-65134, Clone: OKT4) and 5.00 ul CoraLite® Plus 488-conjugated Anti-Human CD25 (CL488-65096, Clone: BC96). Cells were not fixed.



1X10<sup>6</sup> human peripheral blood mononuclear cells (PBMCs) were surface stained with APC Anti-Human CD4 (APC-65134, Clone: OKT4) and CoraLite® Plus 488-conjugated Mouse IgG1 isotype control antibody. Cells were not fixed.