

Nur für Forschungszwecke

BCL6 Monoklonaler Antikörper

Katalog-Nr.:CL488-66340



Allgemeine Informationen

Katalog-Nr.: CL488-66340	GenBank-Zugangsnummer: BC150184	Reinigungsmethode: Protein-G-Reinigung
Größe: 100ul , Konzentration: 2000 µg/ml von604 Nanodrop;	GeneID (NCBI): B-cell CLL/lymphoma 6	CloneNo.: 1E6B1
Wirt: Maus	Vollständiger Name: B-cell CLL/lymphoma 6	Anregungs-/Emissionsmaxima-Wellenlängen: 491 nm / 516 nm
Isotyp: IgG1	Berechnete Masse: 706 aa, 79 kDa	
Immunogen Katalognummer: AG151519		

Anwendungen

Geprüfte Anwendungen:
FC (Intra)

Getestete Reaktivität:
Human

Hintergrundinformationen

BCL6, a zinc finger transcription factor, contains a N-terminal BTB/POZ domain and C-terminal zinc finger DNA-binding motifs and represses transcription of a wide range of target proteins and microRNAs. BCL6 protein has been reported as a master regulator of B lymphocyte development and growth, and altered BCL6 protein expression was implicated in pathogenesis of diverse human hematologic malignancies, especially in the diffuse large B cell lymphoma (DLBCL). BCL6 is required for the development of T follicular helper T cells (TFH), a helper T cell subset required for the formation of mature and productive GCs. BCL6 has also been shown to play important regulatory roles in macrophages. This antibody only recognize human species.

Lagerung

Lagerungsbedingungen:
Bei -20°C lagern. Vor Licht schützen. Nach dem Versand ein Jahr stabil.

Lagerungspuffer:
BS mit 50% Glycerin, 0,05% Proclin300, 0,5% BSA, pH 7,3.

Aliquotieren ist nicht notwendig bei -20°C Lagerung

***** 20ul-Größen enthalten 0.1% BSA**


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Ausgewählte Validierungsdaten



1X10⁶ Ramos cells were intracellularly stained with 0.4 ug CoraLite®488 Anti-Human BCL6 (CL488-66340, Clone:1E6B1) (red), or 0.4 ug Mouse IgG1 Isotype Control (CL488-66360, Clone: T1F8D3F10) (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).