#### Nur für Forschungszwecke

# IL-6 Monoklonaler Antikörper

Katalog-Nr.:CL488-69001



Reinigungsmethode:

Protein-G-Reinigung

Anregungs-/Emissionsmaxima-

CloneNo.:

Wellenlängen: 493 nm / 522 nm

4G3E6

#### Allgemeine Informationen

GenBank-Zugangsnummer: Katalog-Nr.:

CL488-69001 GeneID (NCBI):

100ul , Konzentration: 1000 µg/ml von<sub>Vollständiger</sub> Name:

Nanodrop; interleukin 6 (interferon, beta 2)

Maus Isotyp: lgG1

Immunogen Katalognummer:

HZ-1019

Anwendungen

Geprüfte Anwendungen:

FC (Intra)

Getestete Reaktivität:

Human

### Hintergrundinformationen

Interleukin-6 (IL-6) is an interleukin that acts as both a pro-inflammatory and anti-inflammatory cytokine. IL-6 protein is secreted by a variety of cell types including T cells and macrophages as phosphorylated and variably glycosylated molecule. IL-6 plays an essential role in the final differentiation of B-cells into Ig-secreting cells involved in lymphocyte and monocyte differentiation. It induces myeloma and plasmacytoma growth and induces nerve cells differentiation Acts on B-cells, T-cells, hepatocytes, hematopoietic progenitor cells and cells of the CNS. IL-6 is also considered a myokine, a cytokine produced from muscle, and is elevated in response to muscle  $contraction. \,IL-6 \,has\,been\,shown\,to\,interact\,with\,interleukin-6\,receptor\,and\,gly coprotein\,130.\,Additionally,\,IL-6\,is$ involved in hematopoiesis, bone metabolism, and cancer progression, and has been defined an essential role in directing transition from innate to acquired immunity.

This antibody is used to Block/Neutralize IL6 protein.

Lagerung

Lagerungsbedingungen:

Bei -20°C lagern. Vor Licht schützen.

Lagerungspuffer:

BS mit 50% Glyzerin, 0,05% Proclin300, 0,5% BSA, pH 7,3.

Aliquotieren ist nicht notwendig bei -20°C lagerung

\*\*\* 20ul-Größen enthalten 0.1% BSA

in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Ausgewählte Validierungsdaten



1X10^6 HUVEC cells untreated (dashed lines) or treated with LPS and Brefeldin A (red) were intracellularly stained with 0.5 ug CoraLite® Plus 488 Anti-Human IL-6 (CL488-69001, Clone:4G3E6), or 0.5 ug Control Antibody (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).