

Nur für Forschungszwecke

P16-INK4A Polyklonaler Antikörper

Katalog-Nr.:CL555-10883

Vorgestelltes Produkt



Allgemeine Informationen

Katalog-Nr.: CL555-10883	GenBank-Zugangsnummer: BC021998	Reinigungsmethode: Antigen-Affinitätsreinigung
Größe: 100ul , Konzentration: 1000 µg/ml von1029 Nanodrop;	GeneID (NCBI): 1029	Anregungs-/Emissionsmaxima-Wellenlängen: 557 nm / 570nm
Wirt: Kaninchen	Vollständiger Name: cyclin-dependent kinase inhibitor 2A (melanoma, p16, inhibits CDK4)	
Isotyp: IgG	Berechnete Masse: 16 kDa	
Immunogen Katalognummer: AG1328	Beobachtete Masse: 16-18 kDa	

Anwendungen

Geprüfte Anwendungen:
FC (Intra)

Getestete Reaktivität:
Human

Hintergrundinformationen

CDKN2A generates several transcript variants which differ in their first exons. At least three alternatively-spliced variants encoding distinct proteins were reported. Two of them named p16-INK4 and p14 are sharing 50% identity. The third one -p14(ARF) is entirely unrelated. 10883-1-AP reacts with p16 specifically. P16 plays an essential role in regulating the cell cycle, and mutations in p16 increase the risk of developing various cancers, including melanoma.

Lagerung

Lagerungsbedingungen:
Bei -20°C lagern. Vor Licht schützen.
Lagerungspuffer:
BS mit 50% Glycerin, 0,05% Proclin300, 0,5% BSA, pH 7,3.
Aliquotieren ist nicht notwendig bei -20°C Lagerung

***** 20ul-Größen enthalten 0.1% BSA**


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Ausgewählte Validierungsdaten



1X10⁶ Jurkat cells were intracellularly stained with 0.2 ug CoraLite®555 Anti-Human P16-INK4A (CL555-10883) (red), or 0.2 ug Control Antibody. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).