

Nur für Forschungszwecke

PEX5 Polyklonaler Antikörper

Katalog-Nr.: CL555-12545

Vorgestelltes Produkt



Allgemeine Informationen

Katalog-Nr.: CL555-12545	GenBank-Zugangsnummer: BC010621	Reinigungsmethode: Antigen-affinitätsgereinigt
Größe: 100ul, Konzentration: 1000 µg/ml von5830	GeneID (NCBI): 5830	Anregungs-/Emissionsmaxima-Wellenlängen: 557 nm / 570nm
Nanodrop;	Vollständiger Name: peroxisomal biogenesis factor 5	
Wirt: Kaninchen	Berechnete Masse: 631 aa, 70 kDa	
Isotyp: IgG	Beobachtete Masse: 68-80 kDa	
Immunogen Katalognummer: AG3268		

Anwendungen

Geprüfte Anwendungen:
FC (Intra)

Getestete Reaktivität:
Human, Maus, Ratte

Hintergrundinformationen

The peroxisomal targeting signal type1 (PTS1) receptor, PEX5, is one member of peroxins (PEXs) which are proteins required for peroxisome assembly. PEX5 and PEX7 function as receptors that recognize PTS1- and PTS2- containing proteins, respectively, and PEX5 binds PTS1 through its C-terminal 40-kDa tetratricopeptide repeat domain. It is a predominantly cytoplasmic, partly peroxisomal protein that appears to shuttle between these compartments as it mediates the import of PTS1-containing proteins. PEX5 has been reported to interact with PEX10, PEX12, PEX13, and PEX14. Defects in PEX5 are a cause of Zellweger syndrome (ZWS), which is a lethal peroxisome biogenesis disorder. This antibody recognizes endogenous PEX5, which migrates with an apparent molecular mass of 80 kDa (PMID: 7790377).

Lagerung

Lagerungsbedingungen:
Bei -20°C lagern. Vor Licht schützen.

Lagerungspuffer:
BS mit 50% Glycerin, 0,05% Proclin300, 0,5% BSA, pH 7,3.

Aliquotieren ist nicht notwendig bei -20°C Lagerung

*** 20ul-Größen enthalten 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Ausgewählte Validierungsdaten



1X10⁶ HeLa cells were intracellularly stained with 0.4 ug Coralite®555 Anti-Human PEX5 (CL555-12545) (red), or 0.4 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).