

Nur für Forschungszwecke

CPT1A Polyklonaler Antikörper

Katalog-Nr.:CL555-15184

Vorgestelltes Produkt



Allgemeine Informationen

| | | |
|---|---|---|
| Katalog-Nr.: CL555-15184 | GenBank-Zugangsnummer: BC000185 | Reinigungsmethode: Antigen-Affinitätsreinigung |
| Größe: 100ul , Konzentration: 1000 µg/ml von1374 Nanodrop; | GeneID (NCBI): 1374 | Anregungs-/Emissionsmaxima-Wellenlängen: 557 nm / 570nm |
| Wirt: Kaninchen | Vollständiger Name: carnitine palmitoyltransferase 1A (Liver) | |
| Isotyp: IgG | Berechnete Masse: 88 kDa | |
| Immunogen Katalognummer: AG7202 | Beobachtete Masse: 86 kDa | |

Anwendungen

Geprüfte Anwendungen:
FC (Intra)

Getestete Reaktivität:
Human, Maus, Ratte

Hintergrundinformationen

CPT1A, also named as CPT1, CPT1-L and L-CPTI, belongs to the carnitine/choline acetyltransferase family. It is localized on Chromosome 11q13.1-2. Carnitine palmitoyltransferase (CPT) deficiencies are common disorders of mitochondrial fatty acid oxidation. The CPT system is made up of two separate proteins located in the outer (CPT1) and inner (CPT2) mitochondrial membranes. CPT1A is an active form of related liver-type carnitine palmitoyltransferase I. (PMID: 11001805). CPT1A deficiency presents as recurrent attacks of fasting hypoketotic hypoglycemia. (PMID: 15363638). This antibody can bind the close sequences genes.

Lagerung


Lagerungsbedingungen:
Bei -20°C lagern. Vor Licht schützen. Nach dem Versand ein Jahr stabil.
Lagerungspuffer:
BS mit 50% Glycerin, 0,05% Proclin300, 0,5% BSA, pH 7,3.
Aliquotieren ist nicht notwendig bei -20°C Lagerung

***** 20ul-Größen enthalten 0.1% BSA**

For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Ausgewählte Validierungsdaten



1X10⁶ HeLa cells were intracellularly stained with 0.8 ug Coralite®555 Anti-Human CPT1A (CL555-15184) (red), or 0.8 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).