

Nur für Forschungszwecke

NeutraKine® IFN Gamma Monoklonaler Antikörper

Katalog-Nr.:CL555-69007



Allgemeine Informationen

Katalog-Nr.: CL555-69007	GenBank-Zugangsnummer: GeneID (NCBI): 3458	Reinigungsmethode: Protein-G-Reinigung
Größe: 100ul, Konzentration: 1000 µg/ml von Nanodrop;	Vollständiger Name: IFN gamma	CloneNo.: 1E10G7
Wirt: Maus		Anregungs-/Emissionsmaxima- Wellenlängen: 557 nm / 570nm
Isotyp: IgG1		
Immunogen Katalognummer: HZ-1301		

Anwendungen

Geprüfte Anwendungen:
FC (Intra)

Getestete Reaktivität:
Human

Hintergrundinformationen

Interferon gamma (IFNG) is a soluble cytokine that is the only member of the type II class of interferons. It is secreted by Th1 cells, cytotoxic T cells and NK cells. The cytokine is associated with antiviral, immunoregulatory and anti-tumor properties and is a potent activator of macrophages. It plays crucial roles in pathogen clearance. Aberrant IFNG expression is associated with a number of autoinflammatory and autoimmune diseases. It has been identified in many studies as a biomarker for pleural tuberculosis (TB). Mutations in this gene are associated with aplastic anemia.

This antibody can be used to neutralize the bioactivity of Interferon gamma.

Lagerung

Lagerungsbedingungen:
Bei -20°C lagern. Vor Licht schützen. Nach dem Versand ein Jahr stabil.

Lagerungspuffer:
BS mit 50% Glycerin, 0,05% Proclin300, 0,5% BSA, pH 7,3.

Aliquotieren ist nicht notwendig bei -20°C Lagerung

*** 20ul-Größen enthalten 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Ausgewählte Validierungsdaten



1X10⁶ PMA and ionomycin treated Jurkat cells were intracellularly stained with 0.25 μ g CoraLite[®]555 Anti-Human NeutraKine[®] IFN Gamma (CL555-69007, Clone:1E10G7), and 0.25 μ g α . Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).