

Nur für Forschungszwecke

Cytokeratin 18 Polyklonaler Antikörper

Katalog-Nr.:CL594-10830



Allgemeine Informationen

Katalog-Nr.: CL594-10830	GenBank-Zugangsnummer: BC000180	Reinigungsmethode: Antigen-Affinitätsreinigung
Größe: 100ul , Konzentration: 1000 µg/ml von3875 Nanodrop;	GeneID (NCBI): keratin 18	Empfohlene Verdünnungen: WB 1:1000-1:4000 IF 1:50-1:500
Wirt: Kaninchen	Vollständiger Name: keratin 18	Anregungs-/Emissionsmaxima-Wellenlängen: 588 nm / 604 nm
Isotyp: IgG	Berechnete Masse: 48 kDa	
Immunogen Katalognummer: AG1260	Beobachtete Masse: 48 kDa	

Anwendungen

Geprüfte Anwendungen: FC (Intra), IF, WB	Positivkontrollen:
Getestete Reaktivität: Human, Maus	WB : A431-Zellen, HeLa-Zellen, K-562-Zellen, MCF-7-Zellen
	IF : HepG2-Zellen, HeLa-Zellen

Hintergrundinformationen

Keratins are a large family of proteins that form the intermediate filament cytoskeleton of epithelial cells, which are classified into two major sequence types. Type I keratins are a group of acidic intermediate filament proteins, including K9-K23, and the hair keratins Ha1-Ha8. Type II keratins are the basic or neutral counterparts to the acidic type I keratins, including K1-K8, and the hair keratins, Hb1-Hb6. KRT18, also named as CYK18, PIG46 and K18, is the most commonly found members of the intermediate filament gene family. KRT18 is a type I keratin which is found primarily in non squamous epithelia and is present in a majority of adenocarcinomas and ductal carcinomas but not in squamous cell carcinomas.

Lagerung

Lagerungsbedingungen:
Bei -20°C lagern. Vor Licht schützen.
Lagerungspuffer:
BS mit 50% Glycerin, 0,05% Proclin300, 0,5% BSA, pH 7,3.
Aliquotieren ist nicht notwendig bei -20°C Lagerung

***** 20ul-Größen enthalten 0.1% BSA**


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Ausgewählte Validierungsdaten



Various lysates were subjected to SDS PAGE followed by western blot with CL594-10830 (Cytokeratin 18 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (-20°C Methanol) fixed HepG2 cells using CoraLite®594 Cytokeratin 18 antibody (CL594-10830) at dilution of 1:200.



1X10⁶ HeLa cells were intracellularly stained with 0.4 ug CoraLite®594 Anti-Human Cytokeratin 18 (CL594-10830) (red), or 0.4 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).