

# ACE2 Polyklonaler Antikörper

Katalog-Nr.:CL594-21115

Vorgestelltes Produkt

## Allgemeine Informationen

Katalog-Nr.:	GenBank-Zugangsnummer:	Reinigungsmethode:
CL594-21115	BC048094	Antigen-Affinitätsreinigung
Größe:	GenID (NCBI):	Empfohlene Verdünnungen:
100ul , Konzentration: 1000 µg/ml von 59272		IF 1:50-1:500
Nanodrop;	Vollständiger Name:	Anregungs-/Emissionsmaxima-
	angiotensin I converting enzyme	Wellenlängen:
Wirz:	(peptidyl-dipeptidase A) 2	588 nm / 604 nm
Kaninchen	Berechneté Masse:	
Isotyp:	805 aa, 92 kDa	
IgG	Beobachteté Masse:	
Immunogen Katalognummer:	120 kDa, 92 kDa	
AG15455		

## Anwendungen

Geprüfte Anwendungen:	Positivkontrollen:
IF	IF : Mausnierengewebe,
Getestete Reaktivität:	

Human, Maus, Ratte

## Hintergrundinformationen


ACE2 (Angiotensin-converting enzyme 2), also named as ACEH, is a zinc metalloprotease of the ACE family and a critical regulator of the reninangiotensin system. ACE2 has a more restricted tissue distribution than ACE, being found predominantly in the heart, kidneys, and testes although low levels have been detected in a variety of tissues (PMID:15983030). ACE2 has been shown to be a functional receptor of the human coronaviruses SARS-CoV and SARS-CoV-2 (PMID: 32142651). The expression level and expression pattern of human ACE2 in different tissues might be critical for the susceptibility, symptoms, and outcome of 2019-nCoV/SARS-CoV-2 infection (PMID: 32133153). It can be used as a potential therapeutic target of SARS-CoV-2 (PMID: 32125455). The calculated molecular weight of ACE2 is 92kDa but it migrates to 120kDa due to N-glycosylation (PMID:16166094). Sometimes, several cleaved fragments can also be detected as 75kDa, 50 kDa or 37kDa (PMID: 29561187, 22009550, 30759273). It has 2 isoforms produced by alternative splicing. This antibody is specific to ACE2. The location of ACE2 is membrane and cytoplasm, however it accumulates in the nucleus during the mitosis (PMID: 1730413/ PMID: 18292088).

## Lagerung


Lagerungsbedingungen:  
Bei -20°C lagern. Vor Licht schützen.  
Lagerungspuffer:  
BS mit 50% Glyzerin, 0,05% Proclin300, 0,5% BSA, pH 7,3.  
Aliquotieren ist nicht notwendig bei -20°C Lagerung

\*\*\* 20ul-Größen enthalten 0.1% BSA

## Ausgewählte Validierungsdaten



Immunofluorescent analysis of (4% PFA) fixed mouse kidney tissue using Coralite®594 ACE2 antibody (CL594-21115) at dilution of 1:200.



Immunofluorescent analysis of (4% PFA) fixed mouse kidney tissue using Coralite®594 ACE2 antibody (CL594-21115) at dilution of 1:200.