

Nur für Forschungszwecke

Glucocorticoid receptor Polyklonaler Antikörper



Katalog-Nr.: CL594-24050

Vorgestelltes Produkt

Allgemeine Informationen

| | | |
|--|---|--|
| Katalog-Nr.: CL594-24050 | GenBank-Zugangsnummer: BC015610 | Reinigungsmethode: Antigen-Affinitätsreinigung |
| Größe: 100ul, Konzentration: 1000 µg/ml von 2908 Nanodrop; | GeneID (NCBI): nuclear receptor subfamily 3, group C, member 1 (glucocorticoid receptor) | Empfohlene Verdünnungen: IF 1:50-1:500 |
| Wirt: Kaninchen | Vollständiger Name: nuclear receptor subfamily 3, group C, member 1 (glucocorticoid receptor) | Anregungs-/Emissionsmaxima-Wellenlängen: 588 nm / 604 nm |
| Isotyp: IgG | Berechnete Masse: 86 kDa | |
| Immunogen Katalognummer: AG21146 | Beobachtete Masse: 94-97 kDa | |

Anwendungen

| | |
|---|--|
| Geprüfte Anwendungen: FC (Intra), IF | Positivkontrollen: IF : A549-Zellen, |
| Getestete Reaktivität: Human, Maus, Ratte | |

Hintergrundinformationen

NR3C1 is a receptor for glucocorticoids, which owns a dual mode of action: as a transcription factor that binds to glucocorticoid response elements (GRE) and as a modulator of other transcription factors. It is involved in cell proliferation and differentiation and specifically implicated in newborn birth weight, thus providing a biological mechanism by which NR3C1 expression may influence birth weight [PMID:22810058].

Lagerung

Lagerungsbedingungen:
Bei -20°C lagern. Vor Licht schützen.
Lagerungspuffer:
BS mit 50% Glycerin, 0,05% Proclin300, 0,5% BSA, pH 7,3.
Aliquotieren ist nicht notwendig bei -20°C Lagerung

*** 20ul-Größen enthalten 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Ausgewählte Validierungsdaten



Immunofluorescent analysis of (4% PFA) fixed A549 cells using CoraLite®594 Glucocorticoid receptor antibody (CL594-24050) at dilution of 1:200.



1X10⁶ HeLa cells were intracellularly stained with 0.4 ug CoraLite®594 Anti-Human Glucocorticoid receptor (CL594-24050) (red), or 0.4 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).