

Nur für Forschungszwecke

CD8a Monoklonaler Antikörper

Katalog-Nr.:CL594-65069



Allgemeine Informationen

Katalog-Nr.:

CL594-65069

Größe:

100tests , 5 µl/test

Wirt:

Ratte

Isotyp:

IgG2a, kappa

GenBank-Zugangsnummer:

BC030679

GeneID (NCBI):

12525

Vollständiger Name:

CD8 antigen, alpha chain

Reinigungsmethode:

Affinitätsreinigung

CloneNo.:

53-6.7

Anregungs-/Emissionsmaxima-

Wellenlängen:

588 nm / 604 nm

Anwendungen

Geprüfte Anwendungen:

FC

Getestete Reaktivität:

Maus

Hintergrundinformationen

CD8 is a transmembrane glycoprotein composed of two disulfide-linked chains. It can be present as a homodimer of CD8a or as a heterodimer of CD8a and CD8β (PMID: 3264320; 8253791). CD8 is found on most thymocytes. The majority of class I-restricted T cells express mostly the CD8aβ heterodimer while CD8aa homodimers alone have been found on some gut intraepithelial T cells , on some T cell receptor (TCR) γδ T cells and on NK cells (PMID: 2111591; 1831127; 8420975). CD8 acts as a co-receptor that binds to MHC class-I and participates in cytotoxic T cell activation (PMID: 8499079). During T cell development, CD8 is required for positive selection of CD4-/CD8+ T cells (PMID: 1968084).

Lagerung

Lagerungsbedingungen:

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

Lagerungspuffer:

PBS mit 0,1% Natriumazid und 0,5% BSA, pH 7,3.


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Ausgewählte Validierungsdaten



1X10⁶ mouse splenocytes were surface stained with CoraLite® Plus 647-conjugated Anti-Mouse CD4 (CL647-65104, Clone: GK1.5) and 5.00 ul CoraLite®594-conjugated Anti-Mouse CD8a (CL594-65069, Clone: 53-6.7). Cells were not fixed.



1X10⁶ mouse splenocytes were surface stained with CoraLite® Plus 647-conjugated Anti-Mouse CD4 (CL647-65104, Clone: GK1.5) and CoraLite®594-conjugated Rat IgG2a isotype control antibody. Cells were not fixed.