

Nur für Forschungszwecke

ATP5A1 Monoklonaler Antikörper

Katalog-Nr.:CL594-66037



Allgemeine Informationen

Katalog-Nr.: CL594-66037	GenBank-Zugangsnummer: BC064562	Reinigungsmethode: Protein-A-Reinigung
Größe: 100ul , Konzentration: 1000 µg/ml von498 Nanodrop;	GeneID (NCBI): 1B10H3	CloneNo.: 1B10H3
Wirt: Maus	Vollständiger Name: ATP synthase, H+ transporting, mitochondrial F1 complex, alpha subunit 1, cardiac muscle	Empfohlene Verdünnungen: WB 1:500-1:1000 IF 1:50-1:500
Isotyp: IgG2b	Berechnete Masse: 60 kDa	Anregungs-/Emissionsmaxima- Wellenlängen: 588 nm / 604 nm
Immunogen Katalognummer: AG8119	Beobachtete Masse: 50 kDa	

Anwendungen

Geprüfte Anwendungen: IF, WB	Positivkontrollen: WB : HEK-293-Zellen, HeLa-Zellen, MCF-7-Zellen, NIH/3T3-Zellen
Getestete Reaktivität: Affe, Human, Maus, Ratte	IF : HepG2-Zellen,

Hintergrundinformationen

The ATP5A1 gene encodes the α subunit of mitochondrial ATP synthase which produces ATP from ADP in the presence of a proton gradient across the membrane. The mitochondrial ATP synthase, also known as Complex V or F1FO ATP synthase, is a multi-subunit enzyme complex consisting of two functional domains, the F1-containing the catalytic core and the Fo- containing the membrane proton channel. FO domain has 10 subunits: a, b, c, d, e, f, g, OSCP, A6L, and F6. F1 is composed of subunits α , β , γ , δ , ϵ , and a loosely attached inhibitor protein IF1. Recently defect in ATP5A1 has been linked to the fatal neonatal mitochondrial encephalopathy. ATP5A1 is localized in the mitochondria and anti-ATP5A1 can be used as the loading control for mitochondrial or Complex V proteins. This antibody recognizes the endogenous ATP5A1 protein in lysates from various cell lines and tissues. The predicted MW of ATP5A1 is 60 kDa, while it undergoes the transit peptide cleavage to become a mature form around 50-55 kDa. Several isoforms of ATP5A1 exist due to the alternative splicing.

Lagerung

Lagerungsbedingungen:
Bei -20°C lagern. Vor Licht schützen. Nach dem Versand ein Jahr stabil.
Lagerungspuffer:
BS mit 50% Glycerin, 0,05% Proclin300, 0,5% BSA, pH 7,3.
Aliquotieren ist nicht notwendig bei -20°C Lagerung

***** 20ul-Größen enthalten 0.1% BSA**


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Ausgewählte Validierungsdaten



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using CoraLite®594 ATP5A1 antibody (CL594-66037, Clone: 1B10H3) at dilution of 1:100.



Various lysates were subjected to SDS PAGE followed by western blot with CL594-66037 (ATP5A1 antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.