

Nur für Forschungszwecke

CCAR2 Monoklonaler Antikörper

Katalog-Nr.:CL594-66497

Vorgestelltes Produkt



Allgemeine Informationen

Katalog-Nr.: CL594-66497	GenBank-Zugangsnummer: BC018269	Reinigungsmethode: Protein-A-Reinigung
Größe: 100ul , Konzentration: 1000 µg/ml von57805 Nanodrop;	GeneID (NCBI): 57805	CloneNo.: 2D10F8
Wirt: Maus	Vollständiger Name: KIAA1967	Empfohlene Verdünnungen: IF 1:50-1:500
Isotyp: IgG2a	Berechnete Masse: 923 aa, 103 kDa	Anregungs-/Emissionsmaxima- Wellenlängen: 588 nm / 604 nm
Immunogen Katalognummer: AG18374		

Anwendungen

Geprüfte Anwendungen: FC (Intra), IF	Positivkontrollen: IF : HeLa-Zellen,
Getestete Reaktivität: Hausschwein, Human	

Hintergrundinformationen

Cell cycle and apoptosis regulator protein 2 (CCAR2), previously known as DBC1/KIAA1967, regulates diverse cellular functions including transcription, mRNA splicing, metabolism, and circadian cycle. CCAR2 is a critical regulator of cell death or survival following cellular stresses. CCAR2 is a widely expressed protein involved in the regulation of a variety of transcriptional complexes. High expression of CCAR2 correlates with poor outcomes in many human tumor types such as squamous cell carcinoma (SCC).

Lagerung

Lagerungsbedingungen:
Bei -20°C lagern. Vor Licht schützen. Nach dem Versand ein Jahr stabil.
Lagerungspuffer:
BS mit 50% Glycerin, 0,05% Proclin300, 0,5% BSA, pH 7,3.
Aliquotieren ist nicht notwendig bei -20°C Lagerung

*** 20ul-Größen enthalten 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Ausgewählte Validierungsdaten



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using CoraLite®594-conjugated CCAR2 antibody (CL594-66497, Clone: 2D10F8) at dilution of 1:100.



1X10⁶ HeLa cells were intracellularly stained with 0.2 ug CoraLite®594 Anti-Human CCAR2 (CL594-66497, Clone:2D10F8) (red), or 0.2 ug Mouse IgG2a Isotype Control (CL594-66360-2, Clone: K11A1B2A2) (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).