

# NEUROD1 Monoklonaler Antikörper

Katalog-Nr.:CL594-66691

## Allgemeine Informationen

Katalog-Nr.:	GenBank-Zugangsnummer:	Reinigungsmethode:
CL594-66691	BC009046	Protein-G-Reinigung
Größe:	GenID (NCBI):	CloneNo.:
100ul , Konzentration: 1200 µg/ml von 4760	3E10F1	
Nanodrop;	Vollständiger Name:	Anregungs-/Emissionsmaxima-
	neurogenic differentiation 1	Wellenlängen:
Wirz:	Berechneté Masse:	588 nm / 604 nm
Maus	356 aa, 40 kDa	
Isotyp:	Beobachteté Masse:	
IgG1	40-50 kDa	
Immunogen Katalognummer:		
AG27606		

## Anwendungen

Geprüfte Anwendungen:

FC (Intra)

Getestete Reaktivität:

Human

## Hintergrundinformationen

NeuroD is a member of the basic helix-loop-helix (bHLH) family of transcription factors. The basic helix-loop-helix (bHLH) proteins are transcription factors that are required for several aspects of development, including cell type determination, terminal differentiation and sex determination. Members of the myogenic determination family, MyoD, myf5, myogenin and MRF4, all have bHLH domains. These proteins function by forming heterodimers with E-proteins and binding to the canonical E-box sequence CANNTG. Neuro D is expressed transiently in a subset of neurons in the central and peripheral nervous systems at the time of their terminal differentiation into mature neurons. Moreover, ectopic expression of Neuro D in *Xenopus* embryos induces premature differentiation of neuronal precursors and Neuro D can convert presumptive epidermal cells into neurons. The lack of NeuroD in the brain results in severe defects in development. Human mutations have been linked to a number of types of diabetes including type I diabetes mellitus and maturity-onset diabetes of the young. The calculated molecular weight of NEUROD1 is 39 kDa, but the modified NEUROD1 protein is about 45-50 kDa.

## Lagerung

Lagerungsbedingungen:

Bei -20°C lagern. Vor Licht schützen. Nach dem Versand ein Jahr stabil.


Lagerungspuffer:

BS mit 50% Glycerin, 0,05% Proclin300, 0,5% BSA, pH 7,3.

Aliquotieren ist nicht notwendig bei -20°C Lagerung

\*\*\* 20ul-Größen enthalten 0,1% BSA

## Ausgewählte Validierungsdaten



1X10<sup>6</sup> SH-SY5Y cells were intracellularly stained with 0.4 ug CoralLite®594 Anti-Human NEUROD1 (CL594-66691, Clone:3E10F1) (red), or 0.4 ug Mouse IgG1 Isotype Control (CL594-66360, Clone: T1F8D3F10) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).