

HO-1/HMOX1 Monoklonaler Antikörper

Katalog-Nr.: CL594-66743

Allgemeine Informationen

Katalog-Nr.:	GenBank-Zugangsnummer:	Reinigungsmethode:
CL594-66743	BC001491	Protein-A-Reinigung
Größe:	GenID (NCBI):	CloneNo.:
100ul, Konzentration: 1000 µg/ml von 3162 Nanodrop;	3162	2D10A5
Wirt:	Vollständiger Name:	Empfohlene Verdünnungen:
Maus	heme oxygenase (decycling) 1	IF 1:50-1:500
Isotyp:	Berechneté Masse:	Anregungs-/Emissionsmaxima-
IgG2a	33 kDa	Wellenlängen:
Immunogen Katalognummer:	Beobachteté Masse:	588 nm / 604 nm
AG21296	33 kDa	

Anwendungen

Geprüfte Anwendungen:	Positivkontrollen:
FC (Intra), IF	IF : humanes Lebergewebe,
Getestete Reaktivität:	
Hauschwein, Human, Maus, Ratte	

Hintergrundinformationen


Heme oxygenase (HMOX1) catalyzes the first and rate-limiting step in the degradation of heme to yield equimolar quantities of biliverdin Ixa, carbon monoxide (CO), and iron. It has 3 isoforms: HO-1 is highly inducible, whereas HO-2 and HO-3 are constitutively expressed (PMID:10194478). Heme oxygenase-1 (HO-1) is expressed in many tissues and vascular smooth muscle cells, and endothelial cells (PMID:15451051) and has been identified as an important endogenous protective factor induced in many cell types by various stimulants, such as hemolysis, inflammatory cytokines, oxidative stress, heat shock, heavy metals, and endotoxin (PMID: 11522663). And the full-length HO-1 is very unstable and susceptible to truncation that generates an inactive, soluble form (28 kDa) (James R. Reed, Pharmacology, 535-568).

Lagerung


Lagerungsbedingungen:
Bei -20°C lagern. Vor Licht schützen. Nach dem Versand ein Jahr stabil.
Lagerungspuffer:
BS mit 50% Glyzerin, 0,05% Proclin300, 0,5% BSA, pH 7,3.
Aliquotieren ist nicht notwendig bei -20°C Lagerung

*** 20ul-Größen enthalten 0.1% BSA


Ausgewählte Validierungsdaten



Immunofluorescent analysis of (4% PFA) fixed human liver tissue using Coralite®594-conjugated HO-1/HMOX1 antibody (CL594-66743, Clone: 2D10A5) at dilution of 1:100.



Immunofluorescent analysis of (4% PFA) fixed human liver tissue using Coralite®594-conjugated HO-1/HMOX1 antibody (CL594-66743, Clone: 2D10A5) at dilution of 1:100.



1X10⁶ HeLa cells were intracellularly stained with 0.2 ug CoraLite®594 Anti-Human HO-1/HMOX1 (CL594-66743, Clone:2D10A5) (red), or 0.2 ug Mouse IgG2a Isotype Control (CL594-66360-2, Clone: K11A1B2A2) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).